

Ketamine not linked to PTSD in military trauma setting

October 17 2017



(HealthDay)—Ketamine administration is not associated with post-

traumatic stress disorder (PTSD) in the military trauma setting, according to a study published online Oct. 3 in *Anaesthesia*.

Georges Mion, M.D., from Cochin Hospital in France, and colleagues conducted a retrospective study using data from the French Military Health Service for 274 soldiers who survived a war injury in Afghanistan (2010-2012) to examine the correlation between [ketamine](#) and PTSD.

The researchers found that 36 percent of the soldiers suffered from PTSD and 32 percent had received ketamine. Fifty-five and 20 percent of patients in the PTSD group and the no-PTSD group, respectively, had received ketamine. The median injury severity score was 5 for injured soldiers who received ketamine compared with 3 among the soldiers who did not receive ketamine. Only [acute stress disorder](#) and total number of surgical procedures were correlated independently with development of PTSD in a multivariable analysis.

"In this [retrospective study](#), ketamine administration was not a risk factor for the development of [post-traumatic stress disorder](#) in the military trauma setting," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Ketamine not linked to PTSD in military trauma setting (2017, October 17) retrieved 11 September 2024 from <https://medicalxpress.com/news/2017-10-ketamine-linked-ptsd-military-trauma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.