

Lifestyle changes can close regional obesity gap, study finds

October 6 2017

Lifestyle differences are to blame for regional variation in obesity rates in Scotland, research has found.

Genetic factors cannot completely explain why obesity is more common in some areas and not others, scientists say.

Public health initiatives that focus on postcodes in deprived areas could help to close this gap and tackle health inequalities between regions, researchers say.

Rates of obesity vary between different regions of countries. Until now, the relative impact of genetics and lifestyle on this variability was unclear.

Scientists led by the University of Edinburgh examined health information from 11,000 people from across Scotland, to determine whether <u>genetic factors</u> or lifestyle differences were the cause of regional differences in obesity rates.

They looked at health traits related to obesity, including weight and <u>body</u> <u>mass index</u> (BMI), from people living in different regions.

These figures were analysed together with genetic information from DNA tests and records of lifestyle and socio-economic factors.

When the team compared data between geographical regions, they found



that <u>lifestyle factors</u> such as smoking, alcohol, diet and other measures of deprivation, had the biggest impact on differences in obesity rates.

There are major health inequalities between regions in Scotland. People living in the most deprived areas can expect to live up to seven years less than those living in the most affluent regions.

Researchers say helping people to change their diet, activity levels and behaviour could reduce differences in <u>obesity</u> rates, and so help narrow the health divide between regions.

The research published in the journal *Nature Communications*, was funded by the Medical Research Council. Participants for the study were drawn from Generation Scotland, a research resource with health data available from more than 20,000 volunteers.

Lead researcher Professor Chris Haley, of the MRC Human Genetics Unit at the University of Edinburgh, said: "Our findings reveal that the factors that have the greatest impact on regional <u>obesity rates</u> can be modified. This is good news because it means we can do something about the problem and potentially narrow the <u>health</u> gap between areas that are least and worst affected.

"Our research supports the conclusion that if we are to understand and then reduce the causes of <u>health inequalities</u>, we need to take into account both genetic and lifestyle differences between individuals."

More information: Carmen Amador et al, Regional variation in health is predominantly driven by lifestyle rather than genetics, *Nature Communications* (2017). DOI: 10.1038/s41467-017-00497-5



Provided by University of Edinburgh

Citation: Lifestyle changes can close regional obesity gap, study finds (2017, October 6) retrieved 30 April 2024 from https://medicalxpress.com/news/2017-10-lifestyle-regional-obesity-gap.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.