

Misperception from WHI prevent women from benefitting from hormone therapy

October 11 2017

More than a decade after the Women's Health Initiative (WHI) clinical trials, lingering misperceptions regarding hormone therapy (HT) still prevent many women from getting relief from their menopause symptoms. A new study from the University of Virginia Health System uncovers knowledge gaps of clinicians treating postmenopausal women and identifies need for additional education. The study results will be presented at the 2017 Annual Meeting of The North American Menopause Society (NAMS) in Philadelphia, October 11-14.

Despite the fact that published evidence from the WHI suggests that HT is a relatively safe, viable solution for symptomatic menopausal <u>women</u> under age 60 or within 10 years postmenopause, the number of women being prescribed and using hormones continues to decline. To counteract ongoing skepticism amongst medical professionals regarding the safety of hormones, a new medical education program was shared with 1,087 physicians, nurse practitioners (NPs), and physician assistants (PAs).

Participants were asked a series of questions before and after reviewing the educational program to identify knowledge gaps relative to hormone safety and determine if additional education would help to overcome misperceptions. Among physicians, the average percentage of participants who correctly answered test questions related to outcomes of the WHI trials was 57% pre-activity and 70% post-activity; for NPs, the average percentage answering correctly was 49% pre-activity and 68% post-activity; and for PAs, 52% and 70% respectively.



"We had expected physicians to demonstrate better results in both the pre- and post-activity tests. As a result of the added education there was improvement in some key areas including the newer analysis of the WHI showing different risks depending on age of starting HT. Although our data suggest there is still uncertainty about HT risks, differences between findings for estrogen alone or with progestin, the interpretation of WHI follow-up studies, and the benefits and risks of newer medications for menopause symptoms," says Dr. JoAnn Pinkerton, lead author of the study from the University of Virginia Midlife Health Center and NAMS executive director. "This indicates that future educational interventions are warranted to reinforce appropriate treatment so women at menopause can get the symptom relief they need."

Dr. Pinkerton is available for interviews before the presentation at the Annual Meeting.

Provided by The North American Menopause Society

Citation: Misperception from WHI prevent women from benefitting from hormone therapy (2017, October 11) retrieved 5 May 2024 from https://medicalxpress.com/news/2017-10-misperception-whi-women-benefitting-hormone.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.