

Use of non-vitamin K blood-thinners with certain medications associated with increased risk of major bleeding

October 3 2017

Among patients with nonvalvular atrial fibrillation, concurrent use of certain commonly prescribed medications with non-vitamin K oral anticoagulants was associated with an increased risk of major bleeding, according to a study published by *JAMA*.

Oral anticoagulation has been proven to prevent [ischemic strokes](#) and prolong life for [patients](#) with [atrial fibrillation](#). Non-vitamin K oral anticoagulants (NOACs) are being used more frequently because of their ease of administration and comparative efficacy compared with warfarin in reducing blood clots and major bleeding. However, for patients with atrial fibrillation, NOACs are often prescribed with other medications that share metabolic pathways that may increase major bleeding risk.

Chang-Fu Kuo, M.D., Ph.D., of Chang Gung Memorial Hospital, Taoyuan, Taiwan, and colleagues used data from the Taiwan National Health Insurance database and included 91,330 patients with nonvalvular atrial fibrillation who received at least one NOAC prescription of dabigatran, rivaroxaban, or apixaban and estimated the bleeding risk associated with or without the concurrent use of 12 commonly prescribed medications that share [metabolic pathways](#) with NOACs (atorvastatin; digoxin; verapamil; diltiazem; amiodarone; fluconazole; ketoconazole, itraconazole, voriconazole, or posaconazole; cyclosporine; erythromycin or clarithromycin; dronedarone; rifampin; or phenytoin).

Among the patients in the study, 4,770 major bleeding events occurred. The researchers found that concurrent use of amiodarone, fluconazole, rifampin, and phenytoin with NOACs had a significant increase in incidence rates of major bleeding than NOACs alone. Compared with NOAC use alone, the incidence rate for [major bleeding](#) was significantly lower for concurrent use of atorvastatin, digoxin, and erythromycin or clarithromycin and was not significantly different for concurrent use of verapamil; diltiazem; cyclosporine; ketoconazole, itraconazole, voriconazole, or posaconazole; and dronedarone.

The study notes some limitations, including that bleeding risk and anticoagulant treatment in the Asian population have been recognized to be different from the Western population. Therefore, the external generalizability of these results, particularly to a Western population may be limited.

"Physicians prescribing NOAC medications should consider the potential risks associated with concomitant use of other drugs," the authors write.

More information: *JAMA* (2017). jamanetwork.com/journals/jama/.../1001/jama.2017.13883

Provided by The JAMA Network Journals

Citation: Use of non-vitamin K blood-thinners with certain medications associated with increased risk of major bleeding (2017, October 3) retrieved 26 April 2024 from <https://medicalxpress.com/news/2017-10-non-vitamin-blood-thinners-medications-major.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.