


New perspectives on risk of HIV and hepatitis among injecting drug users

October 24 2017, by Marion Downey



Credit: AI-generated image ([disclaimer](#))

Reviews of the global prevalence of injecting drug use and of interventions to prevent the spread of blood-borne viruses among people who inject drugs paint a worrying picture.

The provision of programs to prevent the spread of HIV and hepatitis

among people who inject drugs (PWID) is inadequate in many countries around the world and presents a critical public health problem, comprehensive reviews by Australian researchers from the National Drug and Alcohol Research Centre at UNSW Sydney have found.

The two reviews of the global prevalence of injecting drug use and of interventions to prevent the spread of blood borne viruses among PWID are published today in leading international journal *The Lancet Global Health*.

The authors estimate that 15.6 million people have recently injected drugs. Of these, 18 percent are living with HIV infection and 52 percent test positive for hepatitis C (HCV) antibody.


Yet despite evidence that needle and syringe programs (NSP) and opioid substitution therapy (OST) reduce HIV and HCV infections, they are still not being implemented in many places, and few people can access them in many countries, the authors found.

Australia is one of only four countries worldwide with high coverage of both NSP and OST – the others are Austria, the Netherlands and Norway.

In Australasia, 1.1 percent of PWID are living with HIV compared with 25 percent of PWID in Eastern Europe, 36 percent in Latin America, 18 percent in Sub Saharan Africa and 19 percent in South Asia.

By contrast, the prevalence of hepatitis C among PWID is more evenly spread – 57 percent of the people who inject drugs in Australia and New Zealand test positive for hepatitis C antibodies, compared with 64 percent in Central Europe, 55 percent in North America and 50 percent in East and South-East Asia.

"Across all countries, a substantial number of people who inject drugs are living with HIV or HCV and are exposed to multiple adverse risk environments that increase health harms," says UNSW's Professor Louisa Degenhardt, lead author of the paper reviewing prevalence of injecting drug use and HIV and hepatitis in this population.



Opioid substitution therapy (OST) per 100 people who inject drugs. Credit: Evan Cunningham/NDARC

The reviews of global prevalence of injecting drug use and of provision of programs to prevent the spread of blood borne viruses are the first to be conducted since 2008, although the results are not directly comparable due to different and more sophisticated data collection

techniques, and better country-specific record keeping.

Evidence of injecting drug use was found in an additional 33 countries compared with the last review – predominantly from Sub Saharan Africa.

Just over half (52 percent) of the countries with evidence of injecting [drug](#) use had needle and syringe programs. Medical treatment to encourage reductions in injecting – opioid substitution therapy – was available in less than half of all countries identified (48 percent).

UNSW's Dr Sarah Larney, lead author of the paper on global coverage of interventions, says: "Coverage of HIV and HCV prevention interventions for PWID remains poor and is likely to be insufficient to effectively prevent HIV and HCV transmission. Scaling up of interventions for PWID remains a crucial priority for halting the HIV and HCV epidemics.

"The presence of interventions alone is not sufficient; the greatest prevention benefits are reported when NSP and OST are implemented in high coverage and in combination," Dr Larney adds.

The three regions worldwide with the highest populations of people who inject drugs, East and South-East Asia, Eastern Europe and North America, all had poor provision of needle syringe programs and opioid substitution therapy.

HIV prevalence in these countries was high, ranging from 9 percent in North America to 25 percent in Eastern Europe. By contrast, only 1 percent of people who inject drugs in Australia and New Zealand are living with HIV.

"Several [countries](#) in these regions have experienced recent HIV

outbreaks as well as persistently high HCV prevalence among PWID," write the authors.

For example, Russia, which has almost 2 million people who inject drugs – nearly 30 percent of whom have HIV and 69 percent of whom have hepatitis C – does not provide OST and has very limited access to NSP, the authors found.

Interactive dashboards containing data from this study are available on the [NDARC website](#).

More information: Sarah Larney et al. Global, regional, and country-level coverage of interventions to prevent and manage HIV and hepatitis C among people who inject drugs: a systematic review, *The Lancet Global Health* (2017). [DOI: 10.1016/S2214-109X\(17\)30373-X](https://doi.org/10.1016/S2214-109X(17)30373-X)

Provided by University of New South Wales

Citation: New perspectives on risk of HIV and hepatitis among injecting drug users (2017, October 24) retrieved 6 May 2024 from <https://medicalxpress.com/news/2017-10-perspectives-hiv-hepatitis-drug-users.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--