


# Scientists discover regional differences among chandelier cells

October 10 2017



A-D, Upper panels show confocal projection images indicating the extent of axonal arbors in RFP(+) hippocampal chandelier cells (h-ChCs) (A, C) and RFP(+) neocortical chandelier cells (nc-ChCs) (B, D). The A-P (A, B) and M-L (C, D) extent of ChC axonal arbors in sagittal and coronal sections, respectively, are shown. Asterisks indicate both ends of axonal arbors. Confocal single optical sections in lower right panels represent enlarged images of boxed areas in upper panels showing axonal varicosities (magenta) aligned along the axon initial segments (green) stained with anti-Ankyrin G antibodies. Lower left panels show individual cartridge structures. E, F, 3-D reconstructions of an h-ChC (E) and an

nc-ChC (F) in sagittal sections. Axonal processes are color-coded according to their depth. Credit: Max Planck Florida Institute for Neuroscience


The brain is composed of distinct regions that differ in their functional roles and cellular architecture. For example, the hippocampus is an area well-known for its involvement in memory and its dysfunction in diseases such as Alzheimer's, while the neocortex is involved in functions such as perception, consciousness, and language. The hippocampus has a single, curved cell layer, while the neocortex has six, stacked layers. At the cellular level, although they share canonical types of inhibitory interneurons (INs) and excitatory principal neurons (PNs), it remains largely unknown to what extent a single type in different brain regions displays similarity in gene expression, axonal shape, connectivity, and developmental origins.

To approach this problem, researchers in the laboratory of Hiroki Taniguchi, Ph.D., Research Group Leader at the Max Planck Florida Institute for Neuroscience (MPFI), took advantage of a unique class of INs called [chandelier cells](#). As their name suggests, these neurons have a distinctive, chandelier-like shape. Most importantly, when their axonal processes stretch out and connect to neighboring [cells](#), they almost always do so at the same location on that neuron - the axon initial segment - and nowhere else. These powerful inhibitory connections control the output of hundreds of neighboring excitatory PNs. These stereotypical features and the fact that chandelier cells are present in both the [hippocampus](#) and the [neocortex](#) make them an ideal model to study regional differences in a single canonical neuronal type.

To visualize different cell types in the brain, researchers often need genetic access to that cell - the ability to express a gene of interest only in, say, chandelier cells, but not neighboring neurons. Years ago,

Taniguchi developed a method for studying chandelier cells in the neocortex, but access to those in the hippocampus has remained elusive. Two postdoctoral researchers in Taniguchi's lab, Yugo Ishino, Ph.D. and Michael Yetman, Ph.D., followed in his footsteps, painstakingly screening molecules until they found one that was reliably expressed in hippocampal chandelier cells - cadherin 6. Luckily, a mouse model already existed that allowed the team to take advantage of this gene expression and use it to compare the two cell populations.

Now with the ability to compare these two populations, the team found that chandelier cells in the hippocampus expand twice the size of axonal arbors in the neocortex and made twice as many connections as their counterparts. Additionally, the hippocampal chandelier cells were born several days earlier during embryonic development than the neocortical ones. Lastly, the team identified a gene, calretinin, in the hippocampal cells which was not expressed in the neocortical cells - suggesting the possibility that these cells exhibit different functional properties as well.



A, B, Schematics showing experimental strategies for the heterotopic

transplantation of nc-ChCs (A) and h-ChCs (B). E17 MGE cells from Nkx2.1-CreER;Ai14 embryos induced at E16 were transplanted into the hippocampus of P1 host animals (A). E16 hippocampal cells from Nkx2.1-CreER;Ai14 embryos induced at E12 were transplanted into the neocortex of P1 host animals (B). C, D, Upper panels show confocal projection images representing the M-L extent of ChC axonal arbors in coronal sections. An h-ChC transplanted into the host neocortex and an nc-ChC transplanted into the host hippocampus are shown in C and D, respectively. Asterisks indicate both ends of axonal arbors. Confocal single optical sections in lower right panels represent enlarged images of boxed areas in upper panels showing axonal varicosities (magenta) apposed to axon initial segments (green) stained with anti-Ankyrin G antibodies. Lower left panels show individual cartridge structures. E, F, 3-D reconstructions of an h-ChC transplanted into the host neocortex (E) and an nc-ChC transplanted into the host hippocampus (F) in coronal sections. Axonal processes are color-coded according to their depth. Credit: Max Planck Florida Institute for Neuroscience

The scientists wondered what factors determined regional differences in these cells' characteristics. Were the traits predetermined by genes that were set at the start of the cell's life? Or did the environment in which they 'grew up' have a stronger influence?

To answer this question of nature versus nurture, the researchers took cells that would grow to become chandelier cells in the hippocampus and transplanted them into the neocortex, and took cells that would grow to become neocortical chandelier cells and transplanted them to the hippocampus. Each ended up taking on the characteristics of its new neighbors, implying that the environment that they grew up in strongly influenced the fate of these cells.

These results, Taniguchi explained, show that exquisite modifications of canonical neuronal types in different brain regions may contribute to

their functional diversification. Future studies should elucidate the molecular and cellular mechanisms by which regional environment controls phenotypic variations of neuronal types.

With the novel genetic access to hippocampal chandelier cells developed in the Taniguchi Lab at MPFI, neuroscientists can begin asking questions about the function of these cells within learning and memory circuits. The ability to manipulate specific genes within these hippocampal chandelier cells may allow for more meticulous studies of several diseases, including epilepsy and schizophrenia, in which these neurons have been implicated.

**More information:** Yugo Ishino et al, Regional Cellular Environment Shapes Phenotypic Variations of Hippocampal and Neocortical Chandelier Cells, *The Journal of Neuroscience* (2017). [DOI: 10.1523/JNEUROSCI.0047-17.2017](https://doi.org/10.1523/JNEUROSCI.0047-17.2017)

Provided by Max Planck Florida Institute for Neuroscience

Citation: Scientists discover regional differences among chandelier cells (2017, October 10) retrieved 24 April 2024 from <https://medicalxpress.com/news/2017-10-scientists-regional-differences-chandelier-cells.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------