


Stress has dramatically different effects on male and female mouse brains

October 10 2017



A group of neurons called CA3 react differently to stress in male and female mice. Credit: Rockefeller University

"Remarkable" is not a word you encounter very often in the scientific literature, where the unadorned description of experiments and their

outcomes is the rule. But the adjective makes a bold appearance in a new report from the Rockefeller University, and with good reason.

Published this week in *Nature Communications*, the paper describes "remarkable" differences in the way the brains of males and females respond to [stress](#). The findings, based on experiments with mice, are doubly notable because they occur in a part of the brain not normally associated with [sex differences](#). They also have implications for the treatment of stress-related illnesses, including mood disorders.

"There is a need to include sex differences in neuropsychiatric research and endocrinology because men and women do respond differently to drugs," says first author Jordan Marrocco, a postdoctoral associate in the neuroendocrinology lab of Bruce S. McEwen, Rockefeller's Alfred E. Mirsky Professor.

That is especially important, notes McEwen, as scientists increasingly seek to tailor new medicines to individual patients. "Broadly speaking," he says, "the pharmaceutical industry has followed a one-size-fits-all philosophy and tested drugs predominantly in males. Some drugs have undergone little testing in women when they go into actual clinical use."

This can create serious problems, he says, citing the sleep drug Ambien, which was discovered to have a much stronger effect on women than men after it reached the market.

A gender divide

To investigate possible sex differences in the brain, the scientists focused on a region of the hippocampus known as CA3, which plays a crucial role in the [stress response](#) and is also involved in memory, the regulation of mood, and information processing. Specifically, they subjected male and [female mice](#) to a stressful task (a six-minute forced

swim) and then used a method called TRAP, developed in the Rockefeller lab of Nathaniel Heintz, to look at how [genes](#) in their CA3 neurons responded.

What they found was, as they write, remarkable.

Many more genes (6,472) were altered by [acute stress](#) in CA3 in females than in males (2,474), when compared to unstressed controls. This large difference suggests, according to the study, that there is a clear genetic component in the response to stress. The scientists propose that genes in the parts of the brain that react to environmental stressors, such as the forced swim, may do so in female mice at a much greater rate than in males.

An additional finding underscores just how huge the differences in stress responses are between genders. The scientists identified 1,842 genes affected by stress in both sexes, however the vast majority of these "overlapping" genes responded in opposite ways in females and males—genes that were activated by stress in males were suppressed by stress in females, and vice versa.

Stress-induced diseases

McEwen's group also experimented with mice engineered to carry a variant of the BDNF gene that, in humans, is known to increase the risk of developing stress-induced neuropsychiatric disorders. The researchers subjected these mice to cognitive tests—a common way to determine the impact of stress—and found that females with the gene variant had impaired spatial memory, even without being stressed. Males with the mutation, also unstressed, did not show such deficits.

The findings suggest that ovarian hormones may interact with the mutated gene in a way that increases the females' stress-related memory

impairment. Marrocco is investigating that possibility in a follow-up study.

The research in BDNF-impaired mice also sheds new light on the intricate relationship between genes and the environment, which can have the same effects on an organism through completely different mechanisms. "What we show," Marrocco says, "is that gene expression in the brain of an unstressed mouse carrying this genetic variant is similar to that in a normal mouse who experiences acute stress."

He adds that more research is needed into the biological differences known to exist between male and female stress responses. By identifying the genetic basis of these differences, and beginning to determine their location in the brain, McEwen and his team have taken a critical step in this direction.

More information: Jordan Marrocco et al. A sexually dimorphic pre-stressed translational signature in CA3 pyramidal neurons of BDNF Val66Met mice, *Nature Communications* (2017). [DOI: 10.1038/s41467-017-01014-4](https://doi.org/10.1038/s41467-017-01014-4)

Provided by Rockefeller University

Citation: Stress has dramatically different effects on male and female mouse brains (2017, October 10) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-10-stress-effects-male-female-mouse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.