


# A stronger twist to cytotoxic amyloid fibrils

October 24 2017



Amyloid fibrils. Credit: Universiteit van Amsterdam (UVA)

Researchers from Amsterdam and Enschede have for the first time performed a structural comparison of two types of amyloid fibrils that have been associated with Parkinson's disease. Using a combination of experimental methods they show that a cytotoxic C-terminal truncated form of the alpha-synuclein protein that is abundant in vivo, aggregates into more strongly twisted fibrils that are more exposed to water. The results have been published in the *Journal of the American Chemical Society*.

The research was carried out by PhD students Steven Roeters (Van 't Hoff Institute for Molecular Sciences, University of Amsterdam, under the supervision of professor Sander Woutersen) and Aditya Iyer (AMOLF, under the supervision of professor Vinod Subramaniam, Vrije

Universiteit Amsterdam), in cooperation with Vladimir Kogan and professor Mireille Claessens from the University of Twente (UT).

Using a combination of techniques, among which [atomic force microscopy](#), UV-circular dichroism, X-ray diffraction and [two-dimensional infrared spectroscopy](#), the researchers performed a structural analysis to explain the differences in the aggregation of wild-type alpha-synuclein (WT- $\alpha$ S) and the so-called C-terminal truncated form of the protein lacking 32 amino acids at the C-terminal end (1-108- $\alpha$ S). The latter has recently been observed by other groups to be quite common in vivo. It forms aggregates more rapidly and has been associated with a progressive form of Parkinson's disease.

The researchers conclude that fibrils formed by 1-108- $\alpha$ S are more-strongly twisted and that - perhaps due to the stronger twist - their fibril core is more exposed to water. The distance between the hydrogen-bonded protein sheets within the fibril appears to be greater. These differences between 1-108- $\alpha$ S and WT- $\alpha$ S fibrils are even so pronounced that WT- $\alpha$ S monomers do not grow into fibrils seeded from short pieces of 1-108- $\alpha$ S fibrils.

By elucidating the key structural aspects of the two fibrillar species, especially of the more cytotoxic 1-108- $\alpha$ S fibrils, the researchers hope to provide clues contributing to an understanding of molecular mechanisms underlying Parkinson's and other related amyloid diseases.

**More information:** Aditya Iyer et al. C-terminal truncated  $\alpha$ -synuclein fibrils contain strongly twisted  $\beta$ -sheets, *Journal of the American Chemical Society* (2017). [DOI: 10.1021/jacs.7b07403](https://doi.org/10.1021/jacs.7b07403)

Provided by University of Amsterdam

Citation: A stronger twist to cytotoxic amyloid fibrils (2017, October 24) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-10-stronger-cytotoxic-amyloid-fibrils.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.