

## Tau may predict neurological outcome after cardiac arrest

October 18 2017



(HealthDay)—Serum tau is a promising novel biomarker for prediction



of neurological outcome in patients with cardiac arrest, according to a study published online Oct. 5 in the *Annals of Neurology*.

Niklas Mattsson, M.D., Ph.D., from Lund University in Sweden, and colleagues measured the neuronal protein tau in serum at 24, 48, and 72 hours after cardiac arrest in 689 patients to test serum tau as a predictor of neurological outcome.

The researchers found that increased tau was associated with poor outcome at six months after cardiac arrest (median 38.5 ng/L in poor versus 1.5 ng/L in good outcome, for tau at 72 hours). Prediction of poor outcome was improved with tau compared with using clinical information. Tau cutoffs had low false positive rates for good outcome and retained high sensitivity for poor outcome. For example, at 72 hours, tau had a false positive rate of 2 percent with a sensitivity of 66 percent. Tau also had higher accuracy than serum neuron-specific enolase (NSE) and was significantly associated with overall survival during follow-up.

"[Serum tau] may be significantly better than <u>serum</u> NSE, which is recommended in guidelines and currently used in clinical practice in several countries to predict outcome after <u>cardiac arrest</u>," conclude the authors.

## More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2017 HealthDay. All rights reserved.

Citation: Tau may predict neurological outcome after cardiac arrest (2017, October 18) retrieved 5 May 2024 from https://medicalxpress.com/news/2017-10-tau-neurological-outcome-cardiac.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.