


Transplanted hematopoietic stem cells reverse damage caused by neuro-muscular disorder

October 26 2017



In this reconstituted schematic, hematopoietic stem cells (HSC) transplanted in a mouse model of Friedreich's ataxia differentiate into microglial cells (red) and transfer mitochondrial protein (green) to neurons (blue), preventing neurodegeneration. Credit: Stephanie Cherqui, UC San Diego School of Medicine

Researchers at University of California San Diego School of Medicine report that a single infusion of wildtype hematopoietic stem and progenitor cells (HSPCs) into a mouse model of Friedreich's ataxia (FA) measurably halted cellular damage caused by the degenerative disease.

The findings, published online in the October 25 issue of *Science Translational Medicine*, suggest a potential therapeutic approach for a [disease](#) that currently is considered incurable.

Friedreich's ataxia is an inherited, degenerative neuromuscular disorder that initially impairs motor function, such as gait and coordination, but can lead to scoliosis, heart disease, vision loss and diabetes. Cognitive function is not affected. The disease is progressively debilitating, and ultimately requires full-time use of a wheelchair. One in 50,000 Americans has FA.

FA is caused by reduced expression of a mitochondrial protein called frataxin (FXN) due to a two mutated or abnormal copies of the FXN gene. In their study, Stephanie Cherqui, PhD, associate professor in the UC San Diego School of Medicine Department of Pediatrics, and colleagues used a [transgenic mouse model](#) that expresses two mutant human FXN transgenes, and exhibits the resulting progressive neurological degeneration and muscle weakness.

Human hematopoietic stem and [progenitor cells](#) (HSPCs), derived from bone marrow, have become a primary vehicle for efforts to replace or regenerate [cells](#) destroyed by a variety of diseases. Previous research by Cherqui and colleagues had shown that transplanting wildtype or normal mouse HSPCs resulted in long-term kidney, eye and thyroid preservation in a mouse model of cystinosis, another genetic disorder.

In this study, Cherqui's team transplanted wildtype HSPCs into an FA mouse model, reporting that the HSPCs engrafted and soon

differentiated into macrophages in key regions of the mice's brain and spinal cord where they appeared to transfer wildtype FXN into deficient neurons and muscle cells.

"Transplantation of wildtype [mouse](#) HSPCs essentially rescued FA-impacted cells," said Cherqui, "Fratxin expression was restored. Mitochondrial function in the brains of the transgenic mice normalized, as did in the heart. There was also decreased skeletal muscle atrophy."

The scientists note that the [mouse model](#) is not perfect mirror of human FA. Disease progression is somewhat different and the precise pathology in mice is not fully known. However, Cherqui said the findings are encouraging and point toward a potential treatment for a disease that currently has none.

More information: Celine J. Rocca et al. Transplantation of wild-type mouse hematopoietic stem and progenitor cells ameliorates deficits in a mouse model of Friedreich's ataxia, *Science Translational Medicine* (2017). [DOI: 10.1126/scitranslmed.aaj2347](https://doi.org/10.1126/scitranslmed.aaj2347)

Provided by University of California - San Diego

Citation: Transplanted hematopoietic stem cells reverse damage caused by neuro-muscular disorder (2017, October 26) retrieved 18 April 2024 from <https://medicalxpress.com/news/2017-10-transplanted-hematopoietic-stem-cells-reverse.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
