

Women who get frequent UTIs may reduce risk by drinking plenty of water

October 6 2017

Drinking an additional three pints of water a day may keep the urinary tract infection (UTI) away - at least for women who are prone - suggests a study being presented at IDWeek 2017.

The study found [women](#) at risk of UTIs who increased their [water](#) intake by about that much water every day were nearly half as likely to get UTIs as women who did not.

"While doctors have long assumed this is the case and often recommended that women at risk for UTIs increase their fluid intake, it's never really undergone a prospective trial before," said Thomas M. Hooton, MD, lead author of the study and clinical director of the Division of Infectious Diseases, University of Miami School of Medicine. "It's good to know the recommendation is valid, and that drinking water is an easy and safe way to prevent an uncomfortable and annoying infection."

Women are more likely to get UTIs than men in part because the urethra is shorter, meaning it is easier for bacteria to travel from the rectum and vagina to the bladder. Drinking more fluids increases the rate of flushing of bacteria from the bladder and also likely reduces the concentration of bacteria that enter the bladder from the vagina. This reduces the opportunities for bacteria to attach to cells that line the [urinary tract](#), which is necessary to cause an infection, Dr. Hooton said.

The study included 140 healthy premenopausal women who had at least

three UTIs in the last year and reported low daily fluid intake. Half of the women (70) who served as the [control group](#) continued their usual daily fluid intake, while the remainder were told to drink 1.5 liters of water a day (about three 16-ounce glasses) in addition to their usual daily fluid intake. After one year, women in the control group had 3.1 UTIs on average, whereas those in the water group had 1.6 UTIs on average, a 48 percent reduction. As a result, the water group averaged fewer regimens of antibiotics (1.8) than the limited-water group (3.5), a reduction of 47 percent. Reducing the use of antibiotics helps decrease the risk of antibiotic resistance.

Researchers followed the women throughout the year using visits and telephone calls. They documented that over the course of the study, on average women in the water group increased their daily [water intake](#) by 1.15 liters (about 2-1/2 pints) for a total daily fluid intake (including water and other beverages) of 2.8 liters, whereas women in the control group did not increase the amount of water they drank and had a total daily [fluid](#) intake of 1.2 liters.

"If a woman has recurrent UTIs and is looking for a way to reduce her risk, the evidence suggests that if she increases the amount of water she drinks and stays with it, she'll likely benefit," Dr. Hooton said.

Forty to 60 percent of women will develop a UTI during their lifetimes and one in four have a repeat infection, according to the National Institute of Diabetes and Digestive and Kidney Diseases. UTIs lead to more than 10 million doctor visits a year, according to the National Kidney Foundation.

Provided by Infectious Diseases Society of America

Citation: Women who get frequent UTIs may reduce risk by drinking plenty of water (2017,

October 6) retrieved 4 May 2024 from <https://medicalxpress.com/news/2017-10-women-frequent-utis-plenty.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.