

## Adult survivors of childhood cancer are more likely to develop high blood pressure

November 22 2017

People who survived childhood cancer were more than twice as likely as the general population to have high blood pressure (hypertension) as adults.

The study is published in *Cancer Epidemiology, Biomarkers & Prevention*, a journal of the American Association for Cancer Research, by Todd M. Gibson, PhD, assistant faculty member in the Epidemiology/Cancer Control department at St. Jude Children's Research Hospital in Memphis, Tennessee.

Improvements in treatment have dramatically increased survival rates from pediatric cancers, with about 83 percent of children surviving at least five years and many becoming long-term survivors. Today, an estimated 420,000 Americans are adult survivors of childhood <u>cancer</u>.

However, many suffer long-term side effects. "High <u>blood pressure</u> is an important modifiable risk factor that increases risk of heart problems in everyone. Research has shown that high blood pressure can have an even greater negative impact on survivors of childhood cancer who were treated with cardiotoxic therapies such as anthracyclines or chest radiation," Gibson said.

To assess the prevalence of high blood pressure among survivors of childhood cancer, Gibson and colleagues examined 3,016 adults who were part of the St. Jude Lifetime Cohort Study, which provides ongoing medical assessments of childhood cancer survivors to advance



knowledge of their long-term health outcomes. Participants were considered to have high blood pressure if their <u>systolic blood pressure</u> was 140 or greater, their <u>diastolic blood pressure</u> was 90 or greater, or if they had been previously diagnosed with hypertension and were taking antihypertensive medication.

The study showed that the prevalence of hypertension was 2.6 times higher among <u>childhood cancer survivors</u> than expected, based on age-, sex-, race- and body mass index-specific rates in the <u>general population</u>.

The prevalence of hypertension increased over time: At age 30, 13 percent of the survivors had hypertension; at 40, 37 percent had hypertension, and by age 50, more than 70 percent of the survivors had hypertension. Gibson said the prevalence of hypertension in cancer survivors matched rates in the general population of people about a decade older.

Certain groups of survivors were the most likely to have hypertension: men; non-Hispanic blacks, older survivors, and those who were overweight or obese, the study showed.

The study found that exposure to radiotherapy or chemotherapy were not significantly associated with hypertension.

Gibson said the lack of association between high blood pressure and radiotherapy and chemotherapy was surprising, and suggests that the connection between childhood cancer survival and adult hypertension is multifactorial and worthy of future research. In the meantime, he said, clinicians should be mindful that survivors of <u>childhood cancer</u> are more likely than the general public to develop high blood pressure.

"The good news is that, unlike prior cancer therapy, <u>high blood pressure</u> is a modifiable risk factor," Gibson noted. "Research is needed to



identify effective interventions to prevent hypertension in survivors, but our results emphasize the importance of blood pressure surveillance and management."

Gibson said a limitation of the study is that it was based on blood pressure measurements that were taken at a single study visit. A clinical diagnosis of <u>hypertension</u> typically requires measurements taken at multiple intervals, he explained. Also, Gibson added, the St. Jude Lifetime Cohort is a group of cancer survivors who undergo frequent clinical follow-up, so its participants may have benefited from being monitored and may, therefore, be in better health than survivors who have less comprehensive follow-up.

## Provided by American Association for Cancer Research

Citation: Adult survivors of childhood cancer are more likely to develop high blood pressure (2017, November 22) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2017-11-adult-survivors-childhood-cancer-high.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.