

# Sleep apnea may increase risk of developing Alzheimer's disease

November 10 2017

---



Sleep apnea may increase AD risk . Credit: ATS

Obstructive sleep apnea (OSA) may put elderly people at greater risk of developing Alzheimer's disease (AD), according to new research published online in the American Thoracic Society's *American Journal of Respiratory and Critical Care Medicine*.

In "[Obstructive Sleep Apnea Severity Affects Amyloid Burden in Cognitively Normal Elderly: A Longitudinal Study](#)," researchers report that biomarkers for [amyloid beta](#), the plaque-building peptides associated with Alzheimer's disease, increase over time in [elderly adults](#)

with OSA in proportion to OSA severity. Thus, individuals with more apneas per hour had greater accumulation of brain amyloid over time.

According to the authors, AD is a neurodegenerative disorder that afflicts approximately five million older Americans. OSA is even more common, afflicting from 30 to 80 percent of the elderly, depending on how OSA is defined.

"Several studies have suggested that sleep disturbances might contribute to amyloid deposits and accelerate cognitive decline in those at risk for AD," said Ricardo S. Osorio, MD, senior study author and assistant professor of psychiatry at New York University School of Medicine.

"However, so far it has been challenging to verify causality for these associations because OSA and AD share risk factors and commonly coexist."

He added that the purpose of this study was to investigate the associations between OSA severity and changes in AD biomarkers longitudinally, specifically whether amyloid deposits increase over time in healthy elderly participants with OSA.

The study included 208 participants, age 55 to 90, with normal cognition as measured by standardized tests and clinical evaluations. None of the participants was referred by a sleep center, used continuous positive airway pressure (CPAP) to treat sleep apnea, was depressed, or had a medical condition that might affect their brain function. The researchers performed lumbar punctures (LPs) to obtain participants' cerebrospinal fluid (CSF) soluble A $\beta$  levels, and then used positron emission tomography, or PET, to measure A $\beta$  deposits directly in the brain in a subset of participants.

The study found that more than half the participants had OSA, including

36.5 percent with mild OSA and 16.8 percent with moderate to severe OSA. From the total study sample, 104 participated in a two-year longitudinal study that found a correlation between OSA severity and a decrease in CSF A $\beta$ 42 levels over time. The authors said this finding is compatible with an increase in amyloid deposits in the brain; the finding was confirmed in the subset of participants who underwent amyloid PET, which showed an increase in amyloid burden in those with OSA.

Surprisingly, the study did not find that OSA severity predicted cognitive deterioration in these healthy elderly adults. Andrew Varga, MD, PhD, study coauthor and a physician specializing in sleep medicine and neurology at the Icahn School of Medicine at Mount Sinai in New York, said this finding suggests that these changes were occurring in the preclinical stages of AD.

"The relationship between amyloid burden and cognition is probably nonlinear and dependent on additional factors," he added. This study finding may also be attributable to the study's relatively short duration, highly educated participants and use of tests that fail to discern changes in cognitive abilities that are subtle or sleep-dependent, the authors wrote.

The high prevalence of OSA the study found in these cognitively normal elderly participants and the link between OSA and [amyloid](#) burden in these very early stages of AD pathology, the researchers believe, suggest the CPAP, dental appliances, positional therapy and other treatments for sleep apnea could delay cognitive impairment and dementia in many older adults.

"Results from this study, and the growing literature suggesting that OSA, cognitive decline and AD are related, may mean that age tips the known consequences of OSA from sleepiness, cardiovascular, and metabolic dysfunction to brain impairment," Dr. Osorio said. "If this is the case,

then the potential benefit of developing better screening tools to diagnose OSA in the elderly who are often asymptomatic is enormous."

Provided by American Thoracic Society

Citation: Sleep apnea may increase risk of developing Alzheimer's disease (2017, November 10) retrieved 3 May 2024 from

<https://medicalxpress.com/news/2017-11-apnea-alzheimer-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.