


# Chemists discover a surprising effect of a well-known leukemia drug

November 13 2017



Structures of active center of L-Asparaginases *Rh. rubrum* (a) and *E. coli* (b)  
Credit: Vadim Pokrovsky

Researchers from RUDN University and Institute of Biomedical Chemistry of the Russian Academy of Sciences have identified an alternative mechanism for the effective antitumor drug—an enzyme called L-asparaginase. Some isoenzymes of L-asparaginase block the growth of telomeres (regions of repetitive nucleotide sequences at each end of a chromosome) on DNA molecules, and this limits the number of divisions of a cancer cell. This effect is reported in the *Biochemical and Biophysical Research Communications*.

"L-asparaginase [enzyme](#) has been used for the treatment of [acute lymphoblastic leukemia](#) (cancer of the hematopoietic system) in children for more than 35 years. In recent years, data on its effectiveness in NK/T-cell lymphomas (atype of skin cancer) has been published. We performed a meta-analysis exploring these findings in one of our recent articles," said co-author Vadim Pokrovsky, professor of the Department of Biochemistry (RUDN).

The antitumor effect of this enzyme first became known in the 1950s, when scientists noticed that the serum of guinea pig blood with a high content of L-asparaginase stops the development of leukemia in mice. This enzyme is present in a large number of bacteria, fungi, protozoa, and even plants. For medical purposes, it is extracted from Escherichia coli (E. coli). The gene encoding L-asparaginase is inserted into the producer strain so that the enzyme can be obtained to produce commercial batches.

The antitumor effect of the enzyme is associated with the breaking down of asparagine. Asparagine is an amino acid from which proteins are built. When L-asparaginase cleaves asparagine, many cells that are sensitive to the deficiency of this amino acid are killed. Above all, this refers to tumor cells that are unable to synthesize this amino acid independently.

RUDN University scientists, in collaboration with their colleagues from the laboratory of medical biotechnology (V.N. Orekhovich Institute of Biomedical Chemistry) and the laboratory of combined therapy of tumors (N.N. Blokhin Oncology Research Institute) discovered and investigated another, previously unknown effect of L-asparaginase. To do this, they used the strain of L-asparaginase obtained from another microorganism, Rhodospirillum rubrum. Amino acid substitutions were introduced into this strain in order to alter its biological properties.

It turned out that L-asparaginase can also inhibit (block) telomerase—an enzyme responsible for elongation of terminal DNA regions (telomeres), the length of which determines the possible number of cell divisions. Thus, scientists have discovered that L-asparaginase is capable of limiting the number of divisions of a cancer cell.

The work was divided into several steps. First, scientists isolated L-asparaginase and demonstrated its ability to inhibit telomerase in cancer [cells](#). At the second stage, the same effect was reproduced in vivo. Researchers took human tumor cell cultures and transplanted them to athymic mice (mice with absence of T-cell immunity, which prevents human tumors from growing in mice). The desired effect was reproduced in model organisms—tumor growth was suspended, and this effect correlated with the shortening of telomeres and telomerase inhibition.

The researchers hope to obtain new enzymes with a stronger antitumor effect, which can be realized due to the presence of this dual mechanism. Along with the shortening of telomeres, L-asparaginase might possess other effects that are not yet known to science.

"The search for alternative effects of L-asparaginase, additional potential targets of this and other enzymes would allow us to design new enzyme-based drugs. That is, one enzyme will act immediately on a few metabolic targets of the tumor cell," Pokrovsky concluded.

**More information:** Dmitry D. Zhdanov et al, Rhodospirillum rubrum L-asparaginase targets tumor growth by a dual mechanism involving telomerase inhibition, *Biochemical and Biophysical Research Communications* (2017). [DOI: 10.1016/j.bbrc.2017.08.078](https://doi.org/10.1016/j.bbrc.2017.08.078)

Provided by RUDN University

Citation: Chemists discover a surprising effect of a well-known leukemia drug (2017, November 13) retrieved 16 May 2024 from

<https://medicalxpress.com/news/2017-11-chemists-effect-well-known-leukemia-drug.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.