

Critical link between obesity and diabetes has been identified

November 28 2017



Credit: CC0 Public Domain

UT Southwestern researchers have identified a major mechanism by which obesity causes type 2 diabetes, which is a common complication of being overweight that afflicts more than 30 million Americans and



over 400 million people worldwide.

Researchers found that in obesity, insulin released into the blood by the pancreas is unable to pass through the cells that form the inner lining of <u>blood vessels</u>. As a result, insulin is not delivered to the muscles, where it usually stimulates most of the body's glucose to be metabolized. Blood glucose levels rise, leading to <u>diabetes</u> and its related cardiovascular, kidney and vision problems, said Dr. Philip Shaul, Director of the Center for Pulmonary and Vascular Biology in the Department of Pediatrics at UT Southwestern.

"It was totally unpredicted that a major problem in obesity is the delivery of circulating insulin to your muscle. It was even more surprising that this problem involves immunoglobulins, which are the proteins that make up circulating antibodies," said Dr. Chieko Mineo, Associate Professor of Pediatrics, who is a co-senior author on the report with Dr. Shaul.

The researchers found that obese mice have an unexpected chemical change in their immunoglobulins. "The abnormal immunoglobulins then act on cells lining blood vessels to inhibit an enzyme needed to transfer insulin from the bloodstream into the muscle," said Dr. Shaul, who holds the Associates First Capital Corporation Distinguished Chair in Pediatrics. "Type 2 diabetes patients have the same chemical change, and if we give a mouse immunoglobulins from a type 2 diabetic individual, the mouse becomes diabetic."

The findings reported in *The Journal of Clinical Investigation* may lead to new tools for diabetes risk screening and new avenues for <u>diabetes</u> prevention or treatment. The researchers identified an agent that they could administer to mice that prevents the chemical change in immunoglobulins that occurs with obesity and preserves healthy glucose status. The researchers plan to test this strategy in humans in the near



future.



Dr. Philip Shaul and Dr. Chieko Mineo lead a team that discovered a major mechanism by which obesity causes type 2 diabetes. Credit: UT Southwestern

More information: Keiji Tanigaki et al, Hyposialylated IgG activates endothelial IgG receptor FcγRIIB to promote obesity-induced insulin resistance, *Journal of Clinical Investigation* (2017). DOI: 10.1172/JCI89333

Provided by UT Southwestern Medical Center

Citation: Critical link between obesity and diabetes has been identified (2017, November 28) retrieved 2 May 2024 from https://medicalxpress.com/news/2017-11-critical-link-obesity-diabetes.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.