

Changes in diet may improve life expectancy in Parkinson's patients

November 24 2017



Credit: University of Aberdeen

New research from the University of Aberdeen shows that weight loss in people with Parkinson's disease leads to decreased life expectancy, increased risk of dementia and more dependency on care.

The team, led by Dr Angus Macleod propose that closer monitoring for [weight loss](#) in Parkinson's patients and interventions in those who lose [weight](#), such as a high calorie diet, may improve [life expectancy](#), reduce [dementia](#) and reduce dependence on carers.

The study, published in *Neurology*, followed 275 people with Parkinson's disease and parkinsonian disorders for up to ten years, monitored patients' weight and investigated associations between weight loss and outcomes of the disease. The main findings showed that weight loss is common in Parkinson's disease and in the parkinsonian disorders and can occur in the early stages of the disease. Further analysis showed that this early weight loss is associated with higher risk of becoming dependent (i.e. needing help with activities of daily living), of developing dementia, and of dying.

Although other studies have identified weight loss as a common problem in Parkinson's disease, this is the first to identify the link between weight loss and death, dementia and dependence on carers.

Dr Angus MacLeod who led the study explained: "Weight loss is a common problem in Parkinson's but it wasn't clear before we did this how common it was, mainly because of biases in previous studies, or what the consequences were of weight loss. Our hypothesis was that people who are losing weight were going to have adverse outcomes.

"Our finding that those who lose weight have poorer outcomes is important because reversing weight loss may therefore improve outcomes. Therefore, it is vital that further research investigate whether e.g. high calorie diets will improve outcomes in people with Parkinson's who lose weight.

The study was partially funded by Parkinson's UK. Professor David Dexter, Deputy Director of Research at Parkinson's UK added: "While

other studies have demonstrated that weight loss is common in Parkinson's, this is the first to consider the impact this symptom may have.

"It has yet to be determined whether this quicker progression can be corrected by supplementation with a [high calorie diet](#), however this could be a key potential development."

Provided by University of Aberdeen

Citation: Changes in diet may improve life expectancy in Parkinson's patients (2017, November 24) retrieved 20 March 2024 from <https://medicalxpress.com/news/2017-11-diet-life-parkinson-patients.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|