


# Study finds donor corneas can be safely preserved for longer period

November 10 2017



In this cross section of a cornea in a person with Fuchs' dystrophy, the green rectangles represent endothelial cells. The magenta dots are outgrowths of Descemet's membrane known as guttae. Guttae are bumps on the back surface of the cornea that grow larger as the disease progresses. During Descemet's stripping automated endothelial keratoplasty (DSAEK), Descemet's membrane and the endothelium are stripped away. Only part of the donor cornea is transplanted. This tissue includes a very thin layer of the corneal stroma, and the Descemet's membrane and endothelium. The tissue is held in place by an air bubble until it adheres to the back of the patient's cornea and begins to function, removing excess water from the swollen cornea. Credit: George Rosenwasser, M.D.

Results from a large, national clinical trial show that corneal donor tissue can be safely stored for 11 days without negatively impacting the success of transplantation surgery to restore vision in people with diseases of the cornea. The cornea is the eye's clear outer covering. Currently, donor corneas are generally not used for surgery in the United States if they have been preserved for longer than seven days. Two reports from the Cornea Preservation Time Study (CPTS), which was funded by the National Eye Institute, part of the National Institutes of Health, appear online Nov. 10 in *JAMA Ophthalmology*.

Expanding the window in which [donor](#) tissues can be considered suitable by even just a few days should help safeguard quality donor tissue and access to vision-saving transplantation procedures. The study's lead investigator was Jonathan Lass, M.D., of Case Western Reserve University School of Medicine and University Hospitals Eye Institute, Cleveland. Patients were randomly assigned to one of two treatment groups. The first received corneas preserved up to seven days, and the second received corneas preserved for eight to 14 days.

The U.S. Food and Drug Administration has approved the use of solutions to preserve donated corneas for 14 days. "The current practice of surgeons to use corneas preserved for no longer than seven days is not evidence-based," said Lass, "but rather a practice based on opinion, which hopefully will change with this new evidence."

Researchers looked at three-year graft success rates among a total of 1,090 individuals (1,330 eyes) who underwent transplantation via Descemet's stripping automated endothelial keratoplasty by 70 surgeons at 40 surgical sites. Most of the patients underwent transplantation for Fuchs' [endothelial corneal dystrophy](#), a progressive disease that causes cells to die in the innermost layer of the cornea called the endothelium.

Death of [corneal endothelial cells](#) is a normal part of aging. However, Fuchs' accelerates this cell death. Corneal transplantation is the only treatment available to restore vision.

Overall, the investigators were unable to conclude that three-year success rates were the same for corneas preserved for eight to 14 days compared with up to seven days (92.1 percent versus 95.3 percent). However, parsing out the data, they found that much of the difference between the groups was accounted for by those who received corneas preserved 12 to 14 days. There was no meaningful difference between those who received corneas preserved up to seven days and those who received corneas preserved eight to 11 days.

While it is true that patients who received corneas preserved 12 to 14 days had comparably lower success rates, Dr. Lass emphasized that even their success was impressively high at 89.3 percent, which may support the use of these tissues as logistics dictate.

In a separate analysis of the CPTS data, the investigators analyzed the extent of endothelial cell loss (ECL), that is, the loss of cells in the corneal endothelium that continued to occur as a normal part of aging in the grafted donor cornea three years after it had been transplanted. They looked to see if differences in corneal preservation time affected ECL after three years. They noninvasively measured ECL in the 945 eyes with successful grafts and found that corneas preserved up to seven days had a 37-percent loss of cells versus a 40-percent loss in corneas preserved eight to 14 days. However, taking a closer look at the data revealed that the effect of corneal preservation time on the loss of endothelial cells was comparable from four to 13 days.

Taken together, the separate analyses looking at graft success rates and ECL support the use of corneas stored up to 11 days.

Currently, there is a sufficient supply of [donor corneas](#) available for people in the United States who require transplants. In 2016, nearly 50,000 corneal transplantations were performed and, on average, it took only about three to four weeks for donor tissue from an eye bank to become available for scheduled surgery. However, corneal [donor tissue](#) is scarce elsewhere in the world and corneal disease is the third leading cause of blindness outside the U.S.

"Additionally, factors balancing supply and demand in the U.S. may shift unfavorably as the population with age-related diseases of the cornea grows, and the supply of donors is challenged by the increasing prevalence of conditions in deceased donors that make their [corneas](#) ineligible for donation," said study investigator Marian Macsai, M.D., an ophthalmologist at NorthShore University Health System, Glenview, Illinois, and president of the Cornea Society, a U.S.-based international society of corneal surgeons. Prior to transplantation, tissue from a donor must be rigorously screened for safety and viability. That means ruling out certain infectious and communicable diseases in the donor.

"Finding ideal donor-patient matches requires time, technology and manpower. Lengthening the timeframe for tissue evaluation will ultimately help expand the donor pool," Dr. Macsai said.

Case Western Reserve University and the clinical trial sites plan to distribute this information in an educational campaign targeted to the eye banking and ophthalmology community, and then follow up to see if clinical practice changes.

**More information:** George O. Rosenwasser et al. Effect of Cornea Preservation Time on Success of Descemet Stripping Automated Endothelial Keratoplasty, *JAMA Ophthalmology* (2017). [DOI: 10.1001/jamaophthalmol.2017.4989](https://doi.org/10.1001/jamaophthalmol.2017.4989)

Jonathan H. Lass et al. Corneal Endothelial Cell Loss 3 Years After Successful Descemet Stripping Automated Endothelial Keratoplasty in the Cornea Preservation Time Study, *JAMA Ophthalmology* (2017).  
[DOI: 10.1001/jamaophthalmol.2017.4970](https://doi.org/10.1001/jamaophthalmol.2017.4970)

Provided by National Eye Institute

Citation: Study finds donor corneas can be safely preserved for longer period (2017, November 10) retrieved 26 April 2024 from <https://medicalxpress.com/news/2017-11-donor-corneas-safely-longer-period.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.