

Fish oil component preconditions vision cells to survive future injury or disease

November 30 2017



Credit: Louisiana State University

A team of LSU Health New Orleans scientists discovered that a component of fish oil not only protects cells critical to vision from potentially lethal initial insults, but also from those that occur in the future. The study showed that the omega-3 fatty acid, DHA, and its derivatives "precondition" photoreceptor and retinal pigment epithelial (RPE) cells to survive. The results are published in the November 2017



online issue of the journal Cellular and Molecular Neurobiology.

"Our findings support the proposed concept that DHA and docosanoids (molecules made in the brain at the onset of injury or disease) are responsible for activating sustained cellular mechanisms that elicit longterm <u>preconditioning</u> protection," says Nicolas Bazan, MD, PhD, Boyd Professor and Director of LSU Health New Orleans Neuroscience Center of Excellence.

According to the authors, a preconditioning (PC) stimulus is a sub-lethal or pharmacologic stressor that activates a counter-regulatory protective response to a future lethal stimulus. Preconditioning takes place when, for example, the blood supply to an organ is interrupted for a short time and then reestablished. The <u>protective response</u> from that first injury would carry over to a subsequent blood supply shortage, much like the immunity a vaccine confers against future exposures to disease.

"This happens in the heart, brain and retina, as well as other organs," Dr. Bazan says. "To harness the therapeutic potential of preconditioning, it is very important to identify the molecules directly involved."

Fish oil contains two types of polyunsaturated fatty acids (PUFAs)—omega-3 (docosahexaenoic acid or DHA) and omega-6 (arachidonic acid or AA). They have distinctly different actions. Omega-3 PUFAs and their enzymatic metabolic derivatives, docosanoids, display potent anti-inflammatory and pro-resolving properties in contrast to the pro-inflammatory actions of omega-6 PUFA derivatives. The researchers found that although they are released concomitantly, DHA can alter the action of AA. When they supplemented DHA prior to the oxidative stress insult, the synthesis of protective DHA derivatives increased while AA synthesis decreased over time.



"Our findings demonstrate that DHA and the induction of docosanoid synthesis is necessary for preconditioning protection, and thus daily survival, of photoreceptor and RPE cells," adds Bazan. "Since omega-3 impairments are associated with neuroinflammation, which contributes to photoreceptor cell dysfunction and death, enhancing the synthesis of docosanoids may provide an opportunity for halting or ameliorating debilitating retinal degenerative diseases, such as the dry form of agerelated macular degeneration," concludes Bazan.

More information: Eric J. Knott et al, Retinal Pigment Epithelium and Photoreceptor Preconditioning Protection Requires Docosanoid Signaling, *Cellular and Molecular Neurobiology* (2017). <u>DOI:</u> <u>10.1007/s10571-017-0565-2</u>

Provided by Louisiana State University

Citation: Fish oil component preconditions vision cells to survive future injury or disease (2017, November 30) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2017-11-fish-oil-component-preconditions-vision.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.