

Revisions to guidelines for management and treatment of low back pain bring changes to treatment

November 14 2017



Credit: University of Sydney

If you visit your family doctor with low back pain (LBP), you may be surprised at the treatment options they suggest now.

Recent changes to major international guidelines for the management of LBP mean that general practitioners (GP) are now unlikely to recommend pain medicines which were previously the go-to treatment.

Low back pain is the leading cause of disability worldwide. It is the second most common reason for seeking care from a [family doctor](#). In Australia, low back pain is the number one cause of early retirement and income poverty.

The new guidelines – the UK National Institute for Health and Care Excellence clinical guideline for low-back pain and sciatica, and a [clinical practice guideline](#) from the American College of Physicians – encourage a shift in thinking about the primary care management of [low-back pain](#).

In response to an escalating prescription opioid crisis, and an overwhelming amount of research showing most pain medicines have little to no effect compared to placebo for people with LBP, the guidelines have radically changed their stance on the medicines.

Instead of pain medicines, GPs might suggest non-medicinal approaches including massage, heat, yoga, mindfulness and various types of physiotherapy and psychological therapies.

The results of a University of Sydney review to investigate the current approach and changes to diagnosis and management of LBP were published today in the *Canadian Medical Association Journal*.

"Until now, the recommended approach to help LBP in general practice was to prescribe simple pain medicines such as paracetamol or anti-inflammatories," said lead author Dr Adrian Traeger, researcher from the Musculoskeletal Health Group at the University's School of Public Health.

"These new guidelines suggest avoiding pain medicines initially and discouraging other invasive treatments such as injections and surgery. The recent changes to these guidelines are important and represent a substantial change in thinking on how best to manage LBP – the previous recommendations were in place for decades.

"If you have an uncomplicated case of recent-onset LBP, your doctor may now simply provide advice on how to remain active and non-drug methods for [pain relief](#) such as heat and massage, and arrange to see you in two weeks to make sure the pain has settled.

"If your pain started a long time ago, they might suggest treatments such as yoga, exercise or mindfulness as treatment. Other effective options could include spinal manipulation, acupuncture, or multi-disciplinary rehabilitation programs.

"These revisions to major international guidelines should see changes to practice worldwide.

However Dr Traeger is concerned that without support from Medicare the suggested reforms could place additional financial strain on those suffering from low back pain.

"There will be challenges to providing this type of care. It's currently much easier and cheaper to provide a prescription for an opioid pain [medicine](#) (which is not a long-term solution to [chronic pain](#) and carries a risk of substantial harm) than a course of treatment with a physiotherapist or psychologist," he said.

"Health systems in most industrialised countries, including Australian Medicare, are simply not set up to fund the care that is considered the most appropriate for low back pain right now.

"Without policy changes, it will be difficult for GPs to follow current best practice. However, if Medicare were to make simple changes to improve affordability of alternatives to [pain](#) medicines, not only would it make a GPs job easier, it could result a major impact on the lives of many living with [low back pain](#), including those who rely on opioids. This needs the attention of the Federal Government."

More information: Adrian Traeger et al. Diagnosis and management of low-back pain in primary care, *Canadian Medical Association Journal* (2017). [DOI: 10.1503/cmaj.170527](https://doi.org/10.1503/cmaj.170527)

Provided by University of Sydney

Citation: Revisions to guidelines for management and treatment of low back pain bring changes to treatment (2017, November 14) retrieved 26 April 2024 from <https://medicalxpress.com/news/2017-11-guidelines-treatment-pain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.