

Hair cortisol levels predict which mothers are more likely to suffer postpartum depression

November 14 2017

Researchers from the University of Granada (UGR), who belong to the Brain, Mind and Behavior Research Center (CIMCYC, from its abbreviation in Spanish) and the Faculty of Psychology, have proven that cortisol levels present in the hair of pregnant women during the first or third trimesters of pregnancy may indicate which of them are more likely to suffer postpartum depression.

Their work, published in *PLoS ONE*, showed that hair [cortisol levels](#) in women who developed [postpartum depression](#) were higher throughout pregnancy than those seen in women who hadn't developed it, being that difference statistically more significant during the first and third trimesters.

The UGR researchers carried out follow-up observations of 44 [pregnant women](#) throughout the gestation period and after giving birth. Each trimester, the mothers underwent a series of tests that evaluated their stress and psychopathological symptoms while simultaneously taking hair samples from which the researchers extracted the cortisol corresponding to the last three months.

The following days after labor the researchers evaluated the mothers' emotional state in order to assess who among them had developed postpartum depression.

Additionally, the results of the study showed that the participants who developed postpartum depression showed [higher levels](#) of somatization

during the first trimester. During the second trimester, they showed higher levels of somatization, obsession-compulsion, depression and anxiety, and during the third [trimester](#), they showed higher levels of somatization and pregnancy-specific stress. Therefore, all those symptoms, along with higher levels of cortisol, would be indicators of a future postpartum depression.

As María Isabel Peralta Ramírez, lead researcher of the project, says, the consequences of those results are very important in the prevention of postpartum depression "since they show that there are various altered psychological and hormonal variables throughout the whole gestation period in comparison to those women who will not suffer postpartum depression. Detecting those differences is the key to anticipate the psychological state of the mother as well as the consequences for the baby that said state could mean."

This study's primary goal has been to assess the effects of [psychological stress](#) on the mother throughout the whole gestation period as well as on birth variables, and on the baby's stress and neurodevelopment.

More information: Rafael A. Caparros-Gonzalez et al. Hair cortisol levels, psychological stress and psychopathological symptoms as predictors of postpartum depression, *PLOS ONE* (2017). [DOI: 10.1371/journal.pone.0182817](#)

Provided by University of Granada

Citation: Hair cortisol levels predict which mothers are more likely to suffer postpartum depression (2017, November 14) retrieved 8 April 2024 from <https://medicalxpress.com/news/2017-11-hair-cortisol-mothers-postpartum-depression.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.