


Hip steroid injections associated with bone changes

November 29 2017




A 58-year-old woman with left hip pain. X-ray from one month prior to the steroid/anesthetic injection demonstrates moderate joint space narrowing (arrows) and bony proliferation (arrowheads). Credit: Radiological Society of North America

Osteoarthritis patients who received a steroid injection in the hip had a significantly greater incidence of bone death and collapse compared with control groups, according to new research presented today at the annual meeting of the Radiological Society of North America (RSNA).

Receiving an [injection](#) of a steroid and anesthetic is a common treatment for patients who are experiencing pain and inflammation in a joint, such as the hip, knee or shoulder.

"Changes due to osteoarthritis, such as narrowing in the space between joints and the development of bony proliferations, typically develop slowly over time," said Connie Y. Chang, M.D., radiologist at Massachusetts General Hospital and assistant professor of radiology at Harvard Medical School in Boston. "When reading follow-up radiographs of patients who had received a hip injection, we noticed changes had developed rapidly in some patients."

To determine whether arthritis worsened in patients following a hip steroid/anesthetic injection, Chang and a team of radiologists specializing in musculoskeletal diagnostic imaging and intervention, including hip injections, conducted a study involving 102 patients (age range 19-92, including 62 women) who received X-ray images of the treated hip at the time of the injection and during a follow-up three to nine months later.



X-rays three months after the steroid/anesthetic injection demonstrate complete loss of joint space (arrows), sclerosis of the femoral head (*), and flattening of femoral head articular surface consistent with collapse (arrowheads). Credit: Radiological Society of North America

Two musculoskeletal radiologists independently reviewed the X-ray images of the patients who received the injections and those of two

control groups matched to demographics and follow-up imaging duration. The control groups consisted of 102 patients who had hip X-rays without steroid/anesthetic injection and 44 patients who underwent imaging and a steroid/anesthetic injection in the shoulder.

The radiologists reported new osteonecrosis in 22-24 percent of hip injection patients, compared to 5-9 percent in the hip control group and 5 percent in the shoulder injection control group. They observed bone collapse in the head of the femur bone, located at the top of the femur at the articulation with the pelvic bone, in 15-17 percent of hip injection patients, versus 4 percent of hip control patients and 2 percent of shoulder control patients.

Hip injection patients also showed increased imaging findings of osteoarthritis compared to the control groups, but the differences were not statistically significant.

Dr. Chang noted that patients receiving hip injections have symptoms of hip pain severe enough to require the injection and may be prone to faster progression of bony changes compared to the [control](#) groups. These considerations may be important as some orthopedists are requesting higher steroid doses and injections in younger patients.

"We need to look at what's going on with the steroid/anesthetic injectate and osteoarthritis patients to determine what's causing the changes that occur in some patients," Dr. Chang said. "However, we don't want to deter [patients](#) from getting an injection. These results are enough to warrant an investigation, but not enough to cancel a procedure."

Provided by Radiological Society of North America

Citation: Hip steroid injections associated with bone changes (2017, November 29) retrieved 2

May 2024 from <https://medicalxpress.com/news/2017-11-hip-steroid-bone.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.