

Insomnia linked to alcohol-use frequency among early adolescents, says new psychology study

November 8 2017, by Tom Mclaughlin



Marmorstein says that parents, educators, and therapists should consider insomnia to be a risk marker for alcohol use. Credit: Rutgers University

Insomnia is linked to frequency of alcohol use among early adolescents,

according to new Rutgers University–Camden research.

"Parents, educators, and therapists should consider [insomnia](#) to be a risk marker for alcohol use, and alcohol use a risk marker for insomnia, among early adolescents," writes Rutgers–Camden researcher Naomi Marmorstein in the study, published recently in the journal *Addictive Behaviors*.

Marmorstein, a professor of psychology at Rutgers–Camden, examined the associations between alcohol use and four sleep-related issues: initial insomnia; [daytime sleepiness](#); sleep irregularity, defined as the difference in weekday and weekend bedtimes; and disturbed sleep, characterized as nightmares, snoring, sleepwalking, wetting the bed, and talking in sleep.

When sleep problems were found to be associated with frequency of alcohol use, she examined whether symptoms of [mental health problems](#) or levels of parental monitoring accounted for these associations.

The research focused on seventh- and eighth-grade students participating in the Camden Youth Development Study, an initiative funded by the National Institute on Drug Abuse at the National Institutes of Health. The study examines the development of mental health problems and resilience among at-risk youth.

Youth completed questionnaires in the classroom that asked how long it took for them to fall asleep, what times they usually went to bed on a weekday and on the weekend or vacation night, how often they experienced sleep disturbances, and whether they ever fell asleep in class or had trouble staying awake after school. They were also asked the frequency of any alcohol use in the previous four months.

In addition, students answered questions which were used to assess

depressive symptoms, as well as evidence of conduct disorder symptoms.

Teachers also completed questionnaires, which were analyzed to determine the presence of [attention-deficit hyperactivity disorder](#) symptoms.

Overall, there were associations between alcohol and both insomnia and daytime sleepiness. Importantly, Marmorstein determined that symptoms of mental health problems and [parental monitoring](#) did not account for the link between insomnia and alcohol use.

"These findings indicate that insomnia may be a unique risk marker for alcohol use among young adolescents," she says.

The Rutgers–Camden researcher notes that the findings are consistent with associations found between insomnia and [alcohol](#) among older adolescents and adults.

Provided by Rutgers University

Citation: Insomnia linked to alcohol-use frequency among early adolescents, says new psychology study (2017, November 8) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-11-insomnia-linked-alcohol-use-frequency-early.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
