

Intensive glucose control in type 2 diabetes can have adverse effects

November 15 2017

The common approach of intensive glucose control to achieve low blood sugar targets in type 2 diabetes can increase the risk of mortality, finds a study by Cardiff University.

Looking at routine data from over 300,000 people in the UK, collected between 2004 and 2015, researchers found that lower levels of glycated haemoglobin—typically regarded as being good diabetes [control](#) - were associated with increased [mortality risk](#), compared to moderate levels, especially in conjunction with intensive treatments that could cause hypoglycaemia.

Professor Craig Currie from Cardiff University's School of Medicine said: "Treatment guidelines generally recommend therapeutic strategies that aim for low levels of glucose control, on the understanding that it reduces risk of macrovascular complications such as [coronary artery disease](#) and stroke. Contrary to this belief, our findings show persuasively that there is an association with increased mortality risk and what is considered to be good glucose control, or low HbA1c."

The findings of the study also suggest that neither randomised trials nor observational studies have been able to demonstrate a consistent pattern of association between levels of glucose control and adverse outcome, without any explanation as to why. Thus, the optimal target for glucose control in patients with type 2 diabetes remains uncertain. Furthermore, the pattern of mortality in relation to glucose control differed in relation to differing types of [diabetes drugs](#). Of most concern was an increase in

mortality risk in those with 'good control' with type 2 diabetes who were treated with insulin and other glucose lowering drugs that induce hypoglycaemia.

Professor Currie said: "Serious questions remain about the safety of some glucose lowering drugs, with scientific evidence and opposing opinions being largely ignored."

The study 'The impact of differing [glucose](#)-lowering regimens on the pattern of association between [glucose control](#) and survival' is published in *Diabetes, Obesity and Metabolism*.

More information: Craig J. Currie et al, The impact of differing glucose-lowering regimens on the pattern of association between glucose control and survival, *Diabetes, Obesity and Metabolism* (2017). [DOI: 10.1111/dom.13155](#)

Provided by Cardiff University

Citation: Intensive glucose control in type 2 diabetes can have adverse effects (2017, November 15) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-11-intensive-glucose-diabetes-adverse-effects.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.