

Intranasal ketamine has more minor side effects than intranasal fentanyl in children with acute pain

November 22 2017

Minor adverse events (e.g., bad taste in the mouth and dizziness) occur more frequently with intranasal ketamine than with intranasal fentanyl in children with suspected extremity fractures. That is the primary finding of a study to be published in the December 2017 issue of *Academic Emergency Medicine* (AEM), a journal of the Society for Academic Emergency Medicine (SAEM).

The lead author of the study is Stacy L. Reynolds, MD, director of the Pediatric Emergency Medicine Fellowship in the Department of Emergency Medicine at Carolinas Medical Center/Levine Children's Hospital.

Reynolds and colleagues, in their exploratory analysis, detected no difference in efficacy or [pain relief](#) between the two drugs. Additionally, no patient in either group experienced a serious adverse event, and none of the side effects required intervention.

Based on their findings, Reynolds, et al, recommends a larger, multicenter, noninferiority study to more rigorously evaluate the safety, efficacy, and potential opioid-sparing benefits of intranasal ketamine analgesia for children with acute pain.

"There is less information about ketamine available than is needed to determine its role as first line pain therapy in children. This study starts

to give us the information necessary to determine ketamine's place as a pain medication and its potential to replace opioids," said James R. Miner, MD, FACEP, chief of emergency medicine at the Hennepin County Medical Center and professor of [emergency medicine](#) at the University of Minnesota.

Dr. Miner has done extensive research in acute pain management and sedation, monitoring during resuscitation, and in [health care delivery](#). He is the author of a textbook entitled "Emergency Sedation and Pain Management."

More information: Stacy L. Reynolds et al, Randomized Controlled Feasibility Trial of Intranasal Ketamine Compared to Intranasal Fentanyl for Analgesia in Children with Suspected Extremity Fractures, *Academic Emergency Medicine* (2017). [DOI: 10.1111/acem.13313](https://doi.org/10.1111/acem.13313)

Provided by Society for Academic Emergency Medicine

Citation: Intranasal ketamine has more minor side effects than intranasal fentanyl in children with acute pain (2017, November 22) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-11-intranasal-ketamine-minor-side-effects.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--