

## Magnesium, T2DM link seen in poor-carbohydrate-quality diet

November 1 2017

---



(HealthDay)—The correlation between higher magnesium intake and

reduced risk of type 2 diabetes is stronger in the context of poor-carbohydrate-quality diets, according to a study published online Oct. 4 in *Diabetes Care*.

Adela Hruby, Ph.D., M.P.H., from Tufts University in Boston, and colleagues assessed dietary intake from food frequency questionnaires every four years for participants in the Nurses' Health Study (NHS), the NHS2, and the Health Professionals' Follow-Up Study (69,176, 91,471, and 42,096 participants, respectively). Biennial and supplementary questionnaires were used to ascertain type 2 diabetes status.

Over 28 years of follow-up, the researchers documented 17,130 incident cases of type 2 diabetes. Those with the highest versus the lowest magnesium intake had significantly lower risk of type 2 diabetes in pooled analyses across the three cohorts (pooled multivariate hazard ratio in quintile 5 versus 1: 0.85). The association between higher magnesium intake and lower risk of type 2 diabetes was stronger for participants with [high glycemic index](#) or low cereal fiber than among those with [low glycemic index](#) or high cereal fiber.

"Higher magnesium intake is associated with lower risk of type 2 [diabetes](#), especially in the context of lower-carbohydrate-quality diets," the authors write.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Magnesium, T2DM link seen in poor-carbohydrate-quality diet (2017, November 1)

retrieved 13 July 2024 from <https://medicalxpress.com/news/2017-11-magnesium-t2dm-link-poor-carbohydrate-quality-diet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.