


Pneumonia treatment with vaccines instead of antibiotics

November 21 2017



Credit: AI-generated image ([disclaimer](#))

Mycoplasma bacteria are one of the most common causes of bacterial pneumonia in children. It is still unclear how the disease develops. Researchers from the University Children's Hospital Zurich and UZH have now demonstrated that there are specific immune cells, so-called B cells, which are crucial for recovery from the infection.

The antibodies produced by these cells eliminate mycoplasma in the lungs, whereas these bacteria can persist in the upper respiratory tract. The team of researchers cultured the bacteria with a fluorescent dye and were thus for the first time able to visually track the pathogen in the [lung](#) and upper respiratory tract during an [infection](#). The findings from the newly developed mouse model mimic clinical observations in children, whose upper respiratory tracts still contained mycoplasma following an infection.

Different immune responses in the lung and upper respiratory tract

Infectious disease specialist Patrick Meyer Sauter showed together with the international team that there are significant differences between the lung and the upper respiratory tract in the immune response after an infection. The researchers found more so-called IgM and IgG antibodies in the lung as well as an increase and activation of B cells in local lymph nodes – which meant that pathogens were able to be eliminated within weeks. In contrast, in the upper respiratory tract they found IgA antibodies but no activation of B cells, which corresponds to the observed persistence of the pathogen at this site. In the end, experiments carried out with B cell-deficient mice demonstrated that the transfer of [antibodies](#) into these mice effectively eliminated the bacteria in the lung, but was not able to clear the bacteria in the [upper respiratory tract](#).

"This data is the first proof that the antibody response is crucial to clear pulmonary infections caused by mycoplasma," says Patrick Meyer Sauter. These findings may help develop specific vaccines that elicit antibody responses to protect from infection: "Our work paves the way for the development of vaccines against mycoplasma. Target groups for such vaccines would include young children without treatment alternatives as a result of the emerging resistance to antibiotics against

mycoplasma in certain countries," says Meyer Sauter.

More information: Patrick M Meyer Sauter et al. The role of B cells in carriage and clearance of *Mycoplasma pneumoniae* from the respiratory tract of mice, *The Journal of Infectious Diseases* (2017).

[DOI: 10.1093/infdis/jix559](https://doi.org/10.1093/infdis/jix559)

Provided by University of Zurich

Citation: Pneumonia treatment with vaccines instead of antibiotics (2017, November 21)
retrieved 19 April 2024 from

<https://medicalxpress.com/news/2017-11-pneumonia-treatment-vaccines-antibiotics.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--