

Researchers identify potential autoantigen in aggressive form of kidney disease

November 2 2017

New research reveals the presence of abundant levels of a certain protein in the kidneys of patients with fibrillary glomerulonephritis (FGN). In FGN, large amounts of protein become trapped in the millions of filtering units—or glomeruli—that make up the kidney. The findings, which appear in two different studies in the *Journal of the American Society of Nephrology (JASN)*, suggest that the identified protein may be a potential diagnostic and therapeutic target for FGN.

FGN is an aggressive kidney disease, and nearly half of patients become dependent on dialysis within several years of diagnosis. No unique clinical markers of FGN have been identified to help diagnose FGN, and current tests involve time-consuming analyses with light microscopy, immunofluorescence, and electron microscopy. Once a diagnosis is made, most patients are treated non-specifically with immunosuppressive therapy, which is often not effective and can be associated with significant adverse effects.

Working independently, research teams from the Mayo Clinic in Rochester, Minnesota, and the University of Washington, in Seattle, analyzed the [protein content](#) of glomeruli in patient biopsy specimens. Both groups detected DnaJ [heat shock protein](#) family member B9 (DNAJB9) as an abundant [protein](#) in FGN glomeruli, but not in glomeruli from healthy individuals. An accumulation of antibodies was also found along with DNAJB9 in glomeruli from patients with FGN, suggesting that DNAJB9 and antibodies that are likely directed against the protein contribute to the glomerular deposits that are a hallmark of

FGN. Therefore, DNAJB9 may be an autoantigen, or a normal protein that is the target of an autoimmune response.

"Now that we know that large amounts of this protein are deposited in the kidneys of patients with FGN, future therapies that reduce the rate of deposition of this protein in the kidney and/or dissolve the deposits from the [kidney](#) may prove to be effective in the treatment of this intriguing disease," said Samih Nasr, MD, co-senior author of the Mayo study.

"The discovery of DNAJB9 as a putative autoantigen in FGN paves the way for the development of new diagnostic and therapeutic tools to help care for [patients](#) with this disease," said Kelly Smith, MD PhD, senior author of the University of Washington study.

More information: "DnaJ Heat Shock Protein Family Member B9 Is a Novel Biomarker for Fibrillary GN" [DOI: 10.1681/ASN.2017050566](https://doi.org/10.1681/ASN.2017050566)

"DnaJ Homolog Subfamily B Member 9 Is a Putative Autoantigen in Fibrillary GN," [DOI: 10.1681/ASN.2017050566](https://doi.org/10.1681/ASN.2017050566)

Provided by American Society of Nephrology

Citation: Researchers identify potential autoantigen in aggressive form of kidney disease (2017, November 2) retrieved 8 May 2024 from <https://medicalxpress.com/news/2017-11-potential-autoantigen-aggressive-kidney-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--