

Teenage depression linked to father's depression

November 15 2017



Credit: George Hodan/public domain

Adolescents whose fathers have depressive symptoms are more likely to experience symptoms of depression themselves, finds a new study led by UCL researchers.

While the link between mothers' depression and depression in their children is well-established, the new *Lancet Psychiatry* study is the first

to find an association between depression in fathers and their teenaged children, independent of whether the mother has depression, in a large sample in the general population. The effects of fathers' and mothers' depression on their children's symptoms were similar in magnitude.

"There's a common misconception that mothers are more responsible for their children's mental health, while fathers are less influential - we found that the link between parent and teen depression is not related to gender," said the study's lead author, Dr Gemma Lewis (UCL Psychiatry).

"Family-focused interventions to prevent depression often focus more on mothers, but our findings suggest we should be just as focused on fathers," she said.

The researchers drew on two large longitudinal studies of children: Growing up in Ireland, and the Millennium Cohort Study in England and Wales, using data from 6070 and 7768 families from the two studies, respectively. Parental [depressive symptoms](#) were assessed using a questionnaire when the children were 9 and 7 years old in the two cohorts, and then adolescent depressive symptoms were assessed when the children were 13 and 14 years old. The study samples were population-based, meaning they included people who experienced symptoms of depression but had not sought treatment.

After adjusting for confounding factors such as maternal depression, family income and parental alcohol use, the researchers found that for every 3-point (one standard deviation) increase on the Mood and Feelings Questionnaire (MFQ; a commonly-used measure of depressive symptoms) on the part of fathers, there was an associated 0.2-point increase in the adolescent's MFQ score. The findings were replicated in both independent study samples.

Incidence of depression increases markedly at the beginning of adolescence, so the researchers say that understanding the risk factors at that age can be key to preventing depression later in life.

"Men are less likely to seek treatment for depression. If you're a father who hasn't sought treatment for your depression, it could have an impact on your child. We hope that our findings could encourage men who experience depressive symptoms to speak to their doctor about it," said Dr Lewis.

Previous studies have shown links between [paternal depression](#) and poor behavioural and emotional outcomes in their children, but no large study in the [general population](#) (as opposed to a clinical population) has looked at the link with [adolescent depression](#) while taking into account [maternal depression](#) as well.

"The [mental health](#) of both parents should be a priority for preventing [depression](#) among adolescents. There has been far too much emphasis on mothers but [fathers](#) are important as well," said the study's senior author, Professor Glyn Lewis (UCL Psychiatry).

More information: *Lancet Psychiatry* (2017). [DOI: 10.1016/S2215-0366\(17\)30408-X](#)

Provided by University College London

Citation: Teenage depression linked to father's depression (2017, November 15) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-11-teenage-depression-linked-father.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.