

First transcatheter implant for diastolic heart failure successful

November 15 2017



Dr. Scott Lilly performs a heart catheterization on Janet Wickham at The Ohio State University Wexner Medical Center as part of a study to investigate a new device for diastolic heart failure. Credit: The Ohio State University Wexner

Results presented today at the American Heart Association Scientific Sessions and published in *Circulation* show that a new device designed to treat diastolic heart failure is safe and effective. The first patient in the randomized, blinded study was enrolled at The Ohio State University Ross Heart Hospital, which also enrolled the most patients in the trial.

Diastolic [heart failure](#) (DHF) occurs when the [heart](#) muscle becomes stiff and doesn't relax enough to allow blood to flow from the lungs into the heart. This causes blood to back up in the lower left chamber, then the upper left chamber and into the lungs, causing shortness of breath.

"The therapies we use for people with [systolic heart failure](#) don't work as well for people with DHF. For decades, we've only been able to manage the symptoms of DHF with diuretics," said Dr. Rami Kahwash, an advanced heart [failure](#) cardiologist and director of the Heart and Vascular Research Organization at Ohio State Ross Heart Hospital.

Kahwash and Dr. Scott Lilly, an interventional cardiologist and director of the Structural Heart Disease Program at Ohio State, were part of the Phase II study that evaluated a dime-size implant intended to relieve the high pressure created by blood backing up from the heart into the lungs. During a heart catheterization, doctors inserted the tiny, interatrial shunt device in an opening they created between the upper left and right chambers. This diverts some of the blood from the high pressure left atrial chamber to the low pressure right atrial chamber, much like a detour around a traffic jam.



Next to a dime for perspective, this small wire device is designed to hold open a surgically created hole in the heart of diastolic heart failure patients. Credit: The Ohio State University Wexner Medical Center

The study randomized 44 DHF patients at multiple centers nationwide. Participants received either the treatment device or a sham procedure. After one month, patients were evaluated by comparing pulmonary capillary wedge pressure (PCWP) readings during right heart catheterization exercise. Elevated PCWP readings indicate failure of the left ventricle.

Researchers reported that the interatrial shunt device reduced PCWP readings by 3.5 mmHg, compared to a reduction of 0.5 mmHg in the

control group.

"This device represents a novel therapy for diastolic heart failure, a condition characterized by lifestyle-limiting breathlessness, and one in which traditional heart failure therapies have not proven effective. If the initial experience is substantiated through ongoing clinical trials, this interatrial shunt device may be an opportunity to improve quality of life and exercise capacity for patients that currently have few other options," Lilly said.



Dr. Rami Kahwash examines a patient at The Ohio State University Wexner Medical Center where a study found that a new device designed to treat diastolic heart failure is safe and effective. Credit: The Ohio State University Wexner Medical Center

Risk factors for DHF include high blood pressure, diabetes, obesity and age. Kahwash said it's estimated that within the next five years, DHF will account for more than half of all heart failure cases, and be responsible for more hospital admissions than systolic heart failure.

Ohio State Ross Heart Hospital is already recruiting patients in the next phase of the clinical trials testing the interatrial shunt [device](#), which is made by Corvia Medical Inc..

Provided by The Ohio State University Wexner Medical Center

Citation: First transcatheter implant for diastolic heart failure successful (2017, November 15) retrieved 8 May 2024 from <https://medicalxpress.com/news/2017-11-transcatheter-implant-diastolic-heart-failure.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--