

# Transgender women take triple the number of HIV tests as trans men

November 1 2017, by Marcene Robinson

---



Credit: University at Buffalo

A new University at Buffalo study has shown that HIV testing among transgender adults was higher in those who identified as female, were African-American or Hispanic, or had a history of incarceration.

The preliminary research, which sought to identify HIV testing patterns in urban, [transgender](#) individuals in Western New York, also found that among participants, those with history as a sex worker, who only engaged in sex with males or had access to stable housing were also more

likely to seek HIV testing.

The study was recently published in Volume 2 of *Transgender Health*, the first peer-reviewed journal dedicated to the health care of transgender individuals.

The findings will help health care professionals understand which segments of the transgender population are not receiving HIV testing and tailor interventions to reach them.

"Knowing specific attributes and characteristics linked to testing will assist in better targeting of this often hidden population," says Adrian Juarez, PhD, co-investigator and assistant professor in the UB School of Nursing.

The investigation was also led by Yu-Ping Chang, PhD, Patricia H. and Richard E. Garman Endowed Professor and associate dean for research and scholarship in the UB School of Nursing.

The transgender community experiences numerous barriers to access to medical care, ranging from social discrimination to limited clinical knowledge by care providers of transgender health needs. They also have an increased likelihood of substance abuse, prostitution and other risk behaviors.

Current HIV treatment strategies use seek, [test](#), treat and retain (STTR) models to ensure HIV-negative populations remain negative and positive populations achieve an undetectable viral load to lower the risk of transmission.

While STTR has been successful in several populations, little evidence suggests the intervention is feasible in transgender communities due to the various barriers they face.

To better understand which segments of the transgender community are not receiving testing, the researchers examined survey data from the New York State AIDS Institute Reporting System. The de-identified data, collected from 2007-13, included HIV testing history, perceived HIV risks and related medical care.

They also interviewed 27 self-identified [transgender men](#) and women above the age of 18 who consented to receiving an HIV test at a Western New York community health care organization.

Of the participants, 23 were transgender females, 40 percent were Caucasian and nearly 30 percent African-American, and more than half were current sex workers or had been incarcerated. Nearly all of the respondents reported a history of substance abuse and most were uninsured.

"Substance use disorder (SUD) is a highly prevalent issue in transgender individuals," says Chang. "Research evidence indicated that SUD is significantly associated with HIV and other sexually transmitted diseases. Therefore, SUD should be properly screened and managed."

Participants reported receiving an average of three HIV tests and one STD test before joining the study. The majority of them did not receive an STD test within a year of the study.

The results found that transgender women received, on average, more than three times as many HIV tests as transgender men.

African-American and Hispanic participants also received, on average, more than twice as many HIV tests as Caucasian transgender men and women. The highest average was reported among African-Americans.

Those who had access to stable housing had an average test rate that was

more than twice that of participants who were homeless or lived at a residential drug treatment facility.

Averages were slightly higher among individuals who only had male sex partners, were previously incarcerated or performed sex work.

"Testing rates are likely higher in these populations because of testing availability in an incarcerated setting with an inmate clinic," says Juarez.

"Current evidence also shows that self-perceived HIV risk is a huge motivator for some to seek out an HIV test, which in turn may be the factor as to why trans individuals with only male sex partners and sex workers tested more frequently."

The researchers also found that participants who were enrolled in support groups were more likely to receive a higher number of tests.

They believe an intervention that brings the HIV test to the individual by increasing opportunities at support groups and various forms of housing could increase testing among marginalized populations.

Provided by University at Buffalo

Citation: Transgender women take triple the number of HIV tests as trans men (2017, November 1) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2017-11-transgender-women-triple-hiv-trans.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--