

In vitro fertilization linked with increased risk of congenital heart defects

November 22 2017

A new analysis of published studies found a 45% increased risk of congenital heart defects in newborns when women become pregnant via in vitro fertilization (IVF)/intracytoplasmic sperm injection (ICSI) than through spontaneous conception.

The *Ultrasound in Obstetrics and Gynecology* analysis included 8 studies with 25,856 children obtained from IVF techniques and 287,995 children spontaneously conceived and a total of 2289 [congenital heart defects](#).

Congenital heart defects were found in 1.30% and 0.68% in the IVF/ICSI and spontaneous conception groups, respectively.

"We believe that IVF/ICSI pregnancies present an [increased risk](#) of CHDs as a consequence of early placental dysfunction; however this hypothesis should be demonstrated in future studies. We recommend fetal echocardiography in all pregnancies from IVF/ICSI," said lead author Dr. Paolo Cavorotto, of the IRCCS San Raffaele Hospital, in Milan, Italy.

More information: Veronica Giorgione et al, Congenital heart defects in IVF/ICSI pregnancy: a systematic review and meta-analysis, *Ultrasound in Obstetrics & Gynecology* (2017). [DOI: 10.1002/uog.18932](https://doi.org/10.1002/uog.18932)

Provided by Wiley

Citation: In vitro fertilization linked with increased risk of congenital heart defects (2017, November 22) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-11-vitro-fertilization-linked-congenital-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.