

# Weight reducing diets, predominantly those low in fat, may reduce risk of early death for adults with obesity

November 14 2017

---

Weight reducing diets, mostly low in fat, may reduce risk of early death in adults with obesity, finds a study published in *The BMJ*. However, the researchers were unable to show if there was any effect of weight reducing diets on heart disease and cancer.

The research provides further evidence of the clinical benefits associated with [weight](#) reducing diets, which have already been shown to prevent of type 2 diabetes.

Adults with [obesity](#) have an increased risk of early death, heart disease, some cancers, type 2 diabetes and many other diseases.

Previous research has shown that weight loss programmes can help with obesity and the prevention of type 2 diabetes, however, there is currently limited evidence that dieting can prevent other serious harm for adults with obesity.

The team of researchers based at the University of Aberdeen in Scotland and the University of Auckland in New Zealand reviewed data from randomised controlled trials (RCTs) to assess the effects of weight loss programmes on deaths from all causes as well as from [heart disease](#) and cancer.

Their study involved 54 trials with over 30,000 adults dating from 1966

to 2016 with a minimum follow-up time of one year.

Although not always adequately described, all but one of the trials included weight loss diets with sufficient information to establish that a reduction in fat intake was prescribed. Study design and quality of evidence were taken into account in the interpretation of findings.

The researchers identified high quality evidence from 34 trials that showed that weight reducing diets decrease all cause early death for adults with obesity - an 18% relative reduction in [early death](#), corresponding to six fewer deaths per 1000 participants.

However, there was much less evidence for the effect of weight reducing diets on deaths from heart problems and cancer. The authors point out that there were fewer [trials](#) that reported events for these outcomes, providing lower quality evidence.

They also point out that they were unable to demonstrate differences between adults who had followed either a weight reducing diet alone, a weight reducing diet with exercise advice, or a weight reducing diet which provided an exercise programme to attend.

"We did not have sufficient data to examine whether other types of diet or physical activity influence outcomes or whether certain groups in the population are more or less likely to benefit" say the authors.

Nevertheless, they say "weight reducing diets, usually low in fat and low saturated fat, with or without an exercise component, may reduce premature all-cause mortality in [adults](#) who are obese."

"By implication our data support public health measures to prevent weight gain and facilitate [weight loss](#) using these types of [diet](#)" they conclude.

**More information:** Effects of weight loss interventions for adults who are obese on mortality, cardiovascular disease, and cancer: systematic review and meta-analysis, *BMJ* (2017).

[www.bmj.com/content/359/bmj.j4849](http://www.bmj.com/content/359/bmj.j4849)

Provided by British Medical Journal

Citation: Weight reducing diets, predominantly those low in fat, may reduce risk of early death for adults with obesity (2017, November 14) retrieved 3 May 2024 from

<https://medicalxpress.com/news/2017-11-weight-diets-predominantly-fat-early.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--