


Weight loss through exercise alone does not protect knees

November 28 2017



Baseline and follow-up MRIs of the right knee of two obese women, one who lost no weight (A, B), and one who lost weight (C, D). Credit: Radiological Society of North America

Obese people who lose a substantial amount of weight can significantly slow down the degeneration of their knee cartilage, but only if they lose weight through diet and exercise or diet alone, according to a new MRI


study presented today at the annual meeting of the Radiological Society of North America (RSNA).

Obesity is a major risk factor for osteoarthritis, a [degenerative joint disease](#) that affects more than a third of adults over the age of 60, according to the Centers for Disease Control and Prevention. The knee joint is a common site of osteoarthritis, and in many people the condition progresses until [total knee replacement](#) becomes necessary.

"Once cartilage is lost in osteoarthritis, the disease cannot be reversed," said the study's lead author, Alexandra Gersing, M.D., from the Department of Radiology and Biomedical Imaging at the University of California, San Francisco.

Since cartilage loss cannot be reversed, it is important for people at risk of osteoarthritis or with early signs of the disease to slow the degeneration of cartilage. Weight loss has been shown to slow down cartilage degeneration in overweight and obese individuals, but it was unclear if the method used to lose weight made a difference.

Dr. Gersing and colleagues investigated cartilage degeneration and joint abnormalities over the course of 96 months in overweight and [obese individuals](#) who maintained stable weight and who lost weight via differing regimens.




Knee joint of a patient without weight loss (A) showing severe cartilage defects after 48 months, and intact knee joint of a patient with a substantial amount of weight loss (B). Credit: Radiological Society of North America

The researchers studied 760 men and women with a body mass index of greater than 25 from the Osteoarthritis Initiative, a nationwide research study focused on the prevention and treatment of [knee osteoarthritis](#). The patients either had mild to moderate osteoarthritis or risk factors for the disease. Patients were divided into a group of 380 patients who lost weight, and a control group of 380 patients who lost no weight. The [weight-loss](#) group was further segmented by weight loss method: diet and exercise, diet alone and exercise alone. The researchers used MRI to quantify knee [osteoarthritis](#) at the beginning of the study, at 48 months and at 96 months.

Cartilage degeneration was significantly lower in the weight loss group,

compared to the control group over the 96 months. However, this finding was only present among the patients who lost weight through diet and exercise or diet alone. Although patients who only exercised lost as much weight as patients who dieted alone or dieted and exercised, weight loss through exercise alone showed no significant difference in cartilage degeneration, compared to the group who lost no weight.

"These results add to the hypothesis that solely exercise as a regimen in order to lose weight in overweight and obese adults may not be as beneficial to the knee joint as weight loss regimens involving diet," Dr. Gersing said.



Baseline and follow-up MRIs of the knee of two obese women, one who lost no weight (A, B), and one who lost weight (C, D). Credit: Radiological Society of North America

Provided by Radiological Society of North America

Citation: Weight loss through exercise alone does not protect knees (2017, November 28)
retrieved 24 April 2024 from <https://medicalxpress.com/news/2017-11-weight-loss-knees.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.