

# Bariatric surgery alters liver fatty acid metabolism

December 8 2017

---



(HealthDay)—Bariatric surgery seems to change fatty acid metabolism

in the liver, according to a study published online Nov. 20 in *Diabetes Care*.

Heidi Immonen, M.D., from Turku University Hospital in Finland, and colleagues measured liver fatty acid uptake (LFU), [blood flow](#), and fat content (LFC) in 14 lean individuals (controls) and 25 obese subjects before bariatric surgery and six months afterwards.

The researchers found that increased LFU in obese subjects correlated with body adiposity. Although LFU remained high compared to controls, it did decline postoperatively. LFC was also normalized. Compared to the controls, liver blood flow (per unit volume) was higher in obese subjects at baseline and increased even further postoperatively. As the liver volume decreased, total organ blood flow was unchanged.

"Changes in perfusion may contribute to improved [liver metabolism](#) postoperatively," the authors write.

**More information:** [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Bariatric surgery alters liver fatty acid metabolism (2017, December 8) retrieved 23 April 2024 from <https://medicalxpress.com/news/2017-12-bariatric-surgery-liver-fatty-acid.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--