


# Blood flow altered in brains of preterm newborns vs. full-term infants

December 4 2017



This figure represents the cerebral blood flow (CBF) maps, corresponding anatomical image aligned to the CBF map, and the regions of interest examined. The scale indicates the quantitative value of the CBF map and is expressed in mL/100g/min. The data are from a preterm infant scanned at term age without evidence of brain injury. The insula (see black arrows in panel 'D') may be particularly vulnerable to the added stresses of the preterm infant's life outside the womb. Credit: M. Bouyssi-Kobar, et al., *The Journal of Pediatrics*.

Cerebral blood flow (CBF) of key regions of newborns' brains is altered in very premature infants and may provide an early warning sign of

disturbed brain maturation well before such injury is visible on conventional imaging, according to a prospective, observational study published Dec. 4, 2017 in *The Journal of Pediatrics*.

"During the third trimester of pregnancy, the fetal [brain](#) undergoes an unprecedented growth spurt. To power that growth, [cerebral blood flow](#) increases and delivers the extra oxygen and nutrients needed to nurture normal [brain development](#)," says Catherine Limperopoulos, Ph.D., director of the Developing Brain Research Laboratory at Children's National Health System and senior author of the study. "In full-term pregnancies, these critical brain structures mature inside the protective womb where the fetus can hear the mother and her heartbeat, which stimulates additional brain maturation. For infants born preterm, however, this essential maturation process happens in settings often stripped of such stimuli."

The challenge: How to capture what goes right or wrong in the developing brains of these very fragile newborns? The researchers relied on arterial spin labeling (ASL) magnetic resonance (MR) imaging, a noninvasive technique that labels the water portion of [blood](#) to map how blood flows through infants' brains in order to describe which regions do or do not receive adequate blood supply. The imaging work can be done without a contrast agent since water from arterial blood itself illuminates the path traveled by cerebral blood.

"In our study, very [preterm infants](#) had greater absolute cortical cerebral blood flow compared with full-term infants. Within regions, however, the insula (a region critical to experiencing emotion), anterior cingulate cortex (a region involved in cognitive processes) and auditory cortex (a region involved in processing sound) for preterm infants received a significantly decreased volume of blood, compared with full-term infants. For preterm infants, parenchymal brain injury and the need for cardiac vasopressor support both were correlated with decreased regional

CBF," Limperopoulos adds.

The team studied 98 preterm infants in the study who were born June 2012 to December 2015, were younger than 32 gestational weeks at birth and who weighed less than 1,500 grams. They matched those preemies by gestational age with 104 infants who had been carried to term. The brain MRIs were performed as the infants slept.

Blood flows where it is needed most with areas of the brain that are used more heavily commandeering more oxygen and nutrients. Thus, during brain development, CBF is a good indicator of functional [brain maturation](#) since brain areas that are the most metabolically active need more blood.

"The ongoing maturation of the newborn's brain can be seen in the distribution pattern of cerebral blood flow, with the greatest volume of blood traveling to the brainstem and deep grey matter," says Marine Bouyssi-Kobar, M.S., the study's lead author. "Because of the sharp resolution provided by ASL-MR images, our study finds that in addition to the brainstem and deep grey matter, the insula and the areas of the brain responsible for sensory and motor functions are also among the most oxygenated regions. This underscores the critical importance of these brain regions in early brain development. In preterm infants, the insula may be particularly vulnerable to the added stresses of life outside the womb."

Of note, compromised regional brain structures in adults are implicated in multiple neurodevelopmental disorders. "Altered development of the insula and [anterior cingulate cortex](#) in newborns may represent early warning signs of preterm [infants](#) at greater risk for long-term neurodevelopmental impairments," Limperopoulos says.

**More information:** *Journal of Pediatrics* (2017). [DOI:](#)

[10.1016/j.jpeds.2017.09.083](https://doi.org/10.1016/j.jpeds.2017.09.083)

Provided by Children's National Medical Center

Citation: Blood flow altered in brains of preterm newborns vs. full-term infants (2017, December 4) retrieved 23 April 2024 from

<https://medicalxpress.com/news/2017-12-blood-brains-preterm-newborns-full-term.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.