

Blood pressure control best achieved with a multilevel, multicomponent approach

December 25 2017



Credit: CC0 Public Domain

Current clinical guidelines recommend lower blood pressure targets for the general population, yet control remains elusive for most. A new study suggests that patients with hypertension can best achieve blood pressure

control with a multilevel, multicomponent approach that includes physician- and non-physician-led interventions. The findings from a comparative effectiveness review are published in *Annals of Internal Medicine*.

Hypertension is the leading risk factor for cardiovascular disease. Despite strong evidence that antihypertensive medications and lifestyle modifications reduce [blood pressure](#) and subsequent morbidity and mortality from [cardiovascular disease](#), hypertension control rates hover between 25 and 40 percent, a rate that has remain unchanged for the past 40 years. As such, strategies are needed to increase control rates.

Researchers from Tulane University reviewed published research to assess effectiveness of eight implementation strategies for [blood pressure control](#) in adults with hypertension compared with regular care. They found that multilevel, multicomponent strategies, such as team-based care with medication titration by a nonphysician clinician, team-based care with medication titration by a physician, and multilevel strategies without team-based care, lead to reductions in systolic blood pressure of around 7, 6, and 5 mm Hg, respectively. Patient-level strategies resulted in [systolic blood pressure](#) reductions of around 4 mm Hg for health coaching and 3 mm Hg for home blood pressure monitoring. Trends were similar for diastolic blood pressure reduction.

According to the researchers, these findings provide evidence that multilevel, multicomponent implementation strategies are most useful and should be recommended in clinical practice and public health policy for hypertension control in communities.

More information: Study:

<http://annals.org/aim/article/doi/10.7326/M17-1805>

Editorial: <http://annals.org/aim/article/doi/10.7326/M17-3036>

Provided by American College of Physicians

Citation: Blood pressure control best achieved with a multilevel, multicomponent approach (2017, December 25) retrieved 23 May 2024 from <https://medicalxpress.com/news/2017-12-blood-pressure-multilevel-multicomponent-approach.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.