


Cholera hotspots found at Uganda's borders and lakes

December 28 2017



High risk districts (hotspots) for cholera in Uganda, 2011-2016. Credit: Mohammad Ali, 18 Nov 2017

Uganda is among the countries in sub-Saharan Africa where cholera remains a recurring problem, despite advances in science and technology for prevention, detection and treatment of the infectious disease. Now, researchers reporting in *PLOS Neglected Tropical Diseases* have identified cholera hotspots around Uganda to help target interventions.

Cholera, caused by ingestion of food or water contaminated with the bacterium *Vibrio cholerae* O1 or O139, is estimated to infect up to 4 million people a year worldwide, and lead to tens of thousands of deaths. In sub-Saharan Africa, the fatality rate is more than 2 percent. Epidemics have occurred in Uganda regularly since 1971 and a number of preventive and control measures are currently in place.

In the new work, Mohammad Ali, of the Johns Hopkins Bloomberg School of Public Health, and colleagues studied [district](#) level cholera outbreak data collected by Uganda's Ministry of Health and spanning 2011 through 2016. In addition, existing datasets on population, rainfall, water, sanitation, and hygiene were used. The group identified cholera hotspots as districts whose center fell within a significantly high risk cluster of cases or where a cluster was completely superimposed onto a district.

The group identified 22 districts at high risk of cholera—out of 112 districts total which reported [cholera outbreaks](#) during the study period. 13 of the districts were near the border of the Democratic Republic of Congo (DRC), and 9 were near a border of Kenya. The risk of having

cholera in these hotspots ranged from 2 to 22 times higher than elsewhere in the country. A short distance to the DRC or Kenya border, proximity to internationally shared lakes and river Nile, and higher annual rainfall in a district were all associated with the risk of having cholera in the district.

"The findings of our study could be used as a guide to strengthen the cholera control program in Uganda," the researchers say. "Since a majority of the hotspot districts are near the DRC or Kenya border, it suggests that close collaboration with these countries would be an effective strategy for controlling [cholera](#) in that part of the world."

More information: Bwire G, Ali M, Sack DA, Nakinsige A, Naigaga M, Debes AK, et al. (2017) Identifying cholera "hotspots" in Uganda: An analysis of cholera surveillance data from 2011 to 2016. *PLoS Negl Trop Dis* 11(12): e0006118. doi.org/10.1371/journal.pntd.0006118

Provided by Public Library of Science

Citation: Cholera hotspots found at Uganda's borders and lakes (2017, December 28) retrieved 23 May 2024 from <https://medicalxpress.com/news/2017-12-cholera-hotspots-uganda-borders-lakes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--