

G-quadruplex regulates breast cancerassociated gene

December 21 2017, by Dana Benson



Credit: CC0 Public Domain

For breast cancer, carrying protein CD44s, instead of CD44v, has a survival advantage. Researchers have now discovered a mechanism by which cells can regulate switching between the two proteins, opening options for the development of novel therapeutic strategies to control cancer growth in the future. The study appears in the journal *Genes & Development*.



"In previous studies, we found that switching from CD44v to CD44s is critical for <u>breast cancer</u> progression and metastasis," said corresponding author Dr. Chonghui Cheng, associate professor of molecular and human genetics and of molecular and cellular biology at Baylor College of Medicine. "Here, we studied how cells regulate the switching between the two proteins at the molecular level."

To build CD44 proteins, the genetic information on the DNA is transcribed into RNA and then translated from RNA into a protein. Cells have the choice of translating the information into protein CD44v or CD44s. Breast <u>cancer</u> cells that translate the RNA into protein CD44s have a <u>survival advantage</u>. The mechanism that mediates which protein is produced is called <u>alternative splicing</u>.

"How cancer cells regulate alternative splicing is becoming a fascinating subject of research," said Cheng, who also is at the Lester and Sue Smith Breast Center, part of the National Cancer Institute-designated Dan L Duncan Comprehensive Cancer Center at Baylor College of Medicine. "The consensus is that decisions on which protein should be made rely on specific linear RNA sequences called G-tracts. But emerging evidence suggests that these decisions may also depend on the threedimensional structure of folded linear RNA G-tracts. One example of these three-dimensional structures is G-quadruplex."

G-quadruplex largely regulates switching between CD44v and CD44s contribute

Working with human <u>cells</u> in culture, the researchers asked whether and how G-quadruplex was important for switching between CD44v and CD44s.

"We carried out very defined molecular and biochemical analyses and



provided extensive data that show that G-quadruplex largely regulates the switching between CD44v and CD44s," said co-first author Dr. Jing Zhang, postdoctoral associate in the Cheng lab, whose key contributions were decisive in accomplishing this work. "G-quadruplex structures also are associated with degenerative diseases and aging. If we understand Gquadruplex better, we could be able to provide new insights into how to treat metastatic breast cancer and neurodegenerative diseases and better understand the aging process."

"What has been missing is an appreciation for the role played by folded linear RNA structures such as G-quadruplex in alternative splicing," Cheng said. "If we only look at one-dimensional, linear G-tracts, we might not be able to figure out how splicing is regulated because the key element could be residing within the three-dimensional structure of Gquadruplex, which is the case in this study."

More information: Huilin Huang et al, RNA G-quadruplex secondary structure promotes alternative splicing via the RNA-binding protein hnRNPF, *Genes & Development* (2017). DOI: 10.1101/gad.305862.117

Provided by Baylor College of Medicine

Citation: G-quadruplex regulates breast cancer-associated gene (2017, December 21) retrieved 26 April 2024 from https://medicalxpress.com/news/2017-12-g-quadruplex-breast-cancer-associated-gene.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.