

# New guidelines aim to prevent medics from failing to diagnose patients with liver disease

December 20 2017

---



New guidelines aim to prevent medics from failing to diagnose patients with liver disease. Credit: University of Birmingham

New recommendations, led by experts at the University of Birmingham, have been published to improve the use of liver blood tests.

The recommendations, published in *Gut*, are aimed at helping healthcare workers diagnose patients with [liver disease](#) as well as preventing unnecessary repeat tests for people unlikely to have significant liver disease.

While the number of deaths from other common conditions is falling in the UK, those due to liver disease have been increasing dramatically, with a 400% increase in death rates between 1970 and 2010. Death rates due to liver disease has also risen sharply in younger people, with a 500% increase in the same period for those aged under 65.

Lead author Professor Philip Newsome, Director of the Centre for Liver Research at the University of Birmingham's Institute of Immunology and Immunotherapy, said: "Liver disease develops silently; there may be no signs or symptoms until the complications of liver failure develop.

"Since the current liver [blood](#) tests were developed in the 1950s, they have been the mainstay of liver disease identification. Unfortunately the way liver blood tests are interpreted means that many patients with liver disease are not identified until at a late stage.

"Liver blood or function tests are checked ever more frequently in both primary and secondary care in an attempt to exclude liver disease, for the monitoring of potential adverse effects of drugs on the liver such as statins, and for the investigation of the generally unwell patient.

"These tests often produce an abnormal result, the clinical significance of which is unclear.

"In many cases they are requested in response to non-specific symptoms where there is little potential link between symptoms and likelihood of liver disease, or the blood tests are performed for unrelated reasons such as chronic disease monitoring.

"This frequently leads to a cycle of additional liver blood test testing in an otherwise asymptomatic individual. Notably, many patients referred to hospital with abnormal liver tests do not have any evidence of significant liver disease."

The new guidelines, which cover both adults and children, recommend that initial investigation in patients with potential liver disease should include bilirubin, albumin, ALT, alkaline phosphatase (ALP) and gamma-glutamyltransferase (GGT) plus a complete blood count.

They also include definitions for 'abnormal' blood tests and advise when patients should have liver blood tests, as well as offering advice on pathways and tools for managing patients with abnormal test results.

Professor Newsome adds: "The pathways will be freely disseminated and should be incorporated into primary care IT systems to allow automatic calculation of risk scores when appropriate, to ensure recommendations can be put into practice."

They also state that research is needed to establish the most cost-effective approach to identify patients with alcohol-related [liver disease](#) and non-alcoholic [fatty liver disease](#) at risk of having advanced [liver fibrosis](#).

The recommendations now supersede earlier guidelines in 2000 and were commissioned by the Clinical Services and Standards Committee of the British Society of Gastroenterology.

**More information:** Philip N Newsome et al, Guidelines on the management of abnormal liver blood tests, *Gut* (2017). [DOI: 10.1136/gutjnl-2017-314924](https://doi.org/10.1136/gutjnl-2017-314924)

Provided by University of Birmingham

Citation: New guidelines aim to prevent medics from failing to diagnose patients with liver disease (2017, December 20) retrieved 22 May 2024 from

<https://medicalxpress.com/news/2017-12-guidelines-aim-medics-patients-liver.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.