

Low oxytocin tied to impaired social-emotional function

December 13 2017



(HealthDay)—Low oxytocin levels are related to a lack of emotional

awareness in women with a history of anorexia nervosa (AN), according to a study published recently in the *International Journal of Eating Disorders*.

Cindy Schmelkin, M.D., from Massachusetts General Hospital in Boston, and colleagues assessed fasting serum oxytocin levels in 79 women (19 who were less than 85 percent of ideal body weight [IBW] with AN, 26 who were 90 to 120 percent IBW with a history of AN, and 34 who were 90 to 120 percent IBW with no eating disorder history).

The researchers found that most measures of social-emotional functioning showed impairment in women with AN or a history of AN, versus those with no eating disorder history. In those with a history of AN, oxytocin levels were low, compared to women with no history. Low oxytocin levels were associated with difficulty identifying feelings ($P = 0.008$) and overall alexithymia ($P = 0.0489$) across groups.

"We speculate that low [oxytocin levels](#) may contribute to alexithymia in [women](#) with anorexia nervosa," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Low oxytocin tied to impaired social-emotional function (2017, December 13)
retrieved 26 April 2024 from
<https://medicalxpress.com/news/2017-12-oxytocin-tied-impaired-social-emotional-function.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--