

Survival down with ventricular arrhythmia early post CF-LVAD

December 26 2017



(HealthDay)—For patients post contemporary continuous flow left

ventricular assist device (CF-LVAD) implantation, early ventricular arrhythmias (VAs) are associated with significantly reduced survival, according to a study published online Dec. 20 in *JACC: Clinical Electrophysiology*.

Brian D. Greet, M.D., from the Baylor College of Medicine in Houston, and colleagues examined the incidence, predictors, and associated mortality of pre-implantation, early, and late VAs in 517 patients who underwent CF-LVAD implantation between 2000 and 2015. All VAs were analyzed; those occurring within 30 days of implant were defined as early.

The researchers observed a significant reduction in survival in association with early VAs (hazard ratio, 1.83) compared to late or no VAs. Prior cardiac surgery and pre-CF-LVAD ventricular tachycardia storm were included as pre-implant variables independently predictive of early VAs (odds ratios, 1.9 and 3.15, respectively). From 2000 to 2007, the incidence of early VAs was as high as 47 percent, while from 2008 to 2015, the highest incidence was below 22 percent.

"VAs within thirty days after CF-LVAD implantation are associated with an increased risk of death," the authors write. "Strategies to reduce arrhythmia burden shortly after CF-LVAD implantation warrant further investigation."

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Survival down with ventricular arrhythmia early post CF-LVAD (2017, December 26) retrieved 15 August 2024 from <https://medicalxpress.com/news/2017-12-survival-ventricular-arrhythmia-early-cf-lvad.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.