

Advanced MRI can detect placental perfusion abnormalities in pregnancies complicated by fetal CHD

January 5 2018



Zungho "Wesley" Zun, Ph.D., the study's lead author. Credit: Children's National Health System

In pregnancies complicated by fetal congenital heart disease (CHD),



global placental perfusion was significantly decreased and regional variation of placental perfusion significantly increased as pregnancies progressed, findings that point to non-invasive imaging providing an early warning of placental dysfunction. A Children's National Health System research team is thought to be the first to report non-invasive, whole placenta perfusion imaging in utero in a study published online Nov. 23, 2017 in *Scientific Reports*.

According to the National Heart, Lung and Blood Institute, congenital heart defects are the most common type of birth defect, affecting 8 in 1,000 newborns. Early in pregnancy, the fetal heart follows a parallel developmental course as the placenta, which supplies the growing fetus with oxygen and nutrients while ferrying out waste products. The study authors write that placental arteries are dynamic during pregnancy, remodeling themselves to accommodate increased <u>blood</u> flow as fetuses undergo explosive growth spurts in later stages of pregnancy. If this crucial remodeling does not occur, the placenta may not supply sufficient oxygen and nutrients to the fetus, leading to <u>fetal growth restriction</u> or preeclampsia.

The research team led by Catherine Limperopoulos, Ph.D., enrolled 48 pregnant women who underwent at least one fetal magnetic resonance imaging (MRI) session during their second or third trimester of pregnancy. Thirty-one of the women were healthy volunteers whose mean gestational age was 30 weeks (range: 21 to 39 gestational weeks). Seventeen women were pregnant with fetuses diagnosed with CHD whose mean gestational age was 32 weeks (range: 22 to 38 gestational weeks).

The researchers used velocity-selective arterial spin labeling (VSASL), a powerful MRI technique that directly measures the rate of delivery of arterial blood to organs like the brain. ASL tracks water molecules within the blood as blood flows through arteries, eliminating the need to



use a contrast agent. The team was able to distinguish the placenta <u>perfusion</u> contributions by the fetus and the mother.

"In pregnancies complicated by fetal CHD, global placental perfusion significantly decreased and regional variation of placental perfusion significantly increased with advancing gestational age," says Zungho "Wesley" Zun, Ph.D., the study's lead author.

"Just like the human brain, heart and kidneys—organs that can commandeer heightened blood flow when needed—the placenta may employ an auto-regulatory mechanism to optimize perfusion," adds Limperopoulos, director of Children's Developing Brain Research Laboratory and the study's senior author. "The early increased global placental profusion in pregnancies complicated by CHD may represent an attempt to correct for insufficient fetal <u>blood flow</u>."

The research team writes that the findings demonstrate that placental dysfunction due to CHD can be apparent as early as the second trimester of <u>pregnancy</u> using this imaging technology.

"The predictive value of VSASL imaging, which we continue to study, holds the promise of detecting dysfunction before placental abnormalities become irreversible," Limperopoulos says.

More information: Zungho Zun et al, Non-Invasive Placental Perfusion Imaging in Pregnancies Complicated by Fetal Heart Disease Using Velocity-Selective Arterial Spin Labeled MRI, *Scientific Reports* (2017). DOI: 10.1038/s41598-017-16461-8

Provided by Children's National Medical Center



Citation: Advanced MRI can detect placental perfusion abnormalities in pregnancies complicated by fetal CHD (2018, January 5) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2018-01-advanced-mri-placental-perfusion-abnormalities.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.