

Aversion to holes driven by disgust, not fear, study finds

January 4 2018, by Carol Clark



Clusters of holes, such as those of a lotus seed pod, may be evolutionarily indicative of contamination and disease — visual cues for rotten or moldy food or skin marred by an infection. Credit: Peripitus/Wikipedia Commons

Trypophobia, commonly known as "fear of holes," is linked to a physiological response more associated with disgust than fear, finds a new study published in *PeerJ*.

Trypophobia is not officially recognized in the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders (DSM). Many people, however, report feeling an aversion to clusters of holes—such as those of a honeycomb, a lotus seed pod or even aerated chocolate.

"Some people are so intensely bothered by the sight of these objects that they can't stand to be around them," says Stella Lourenco, a psychologist at Emory University whose lab conducted the study. "The phenomenon, which likely has an evolutionary basis, may be more common than we realize."

Previous research linked tryophobic reactions to some of the same visual spectral properties shared by images of evolutionarily threatening animals, such as snakes and spiders. The repeating pattern of high contrast seen in clusters of holes, for example, is similar to the pattern on the skin of many snakes and the pattern made by a spider's dark legs against a lighter background.

"We're an incredibly visual species," says Vladislav Ayzenberg, a graduate student in the Lourenco lab and lead author of the *PeerJ* study. "Low-level visual properties can convey a lot of meaningful information. These [visual cues](#) allow us to make immediate inferences—whether we see part of a snake in the grass or a whole snake—and react quickly to potential danger."

It is well-established that viewing images of threatening animals generally elicits a fear reaction in viewers, associated with the sympathetic nervous system. The heart and breathing rate goes up and

the pupils dilate. This hyperarousal to potential danger is known as the fight-or-flight response.

The researchers wanted to test whether this same [physiological response](#) was associated with seemingly innocuous images of holes.

They used eye-tracking technology that measured changes in pupil size to differentiate the responses of study subjects to images of clusters of holes, images of threatening animals and neutral images.

Unlike images of snakes and spiders, images of holes elicited greater constriction of the pupils—a response associated with the parasympathetic nervous system and feelings of disgust.

"On the surface, images of threatening animals and clusters of holes both elicit an aversive reaction," Ayzenberg says. "Our findings, however, suggest that the physiological underpinnings for these reactions are different, even though the general aversion may be rooted in shared visual-spectral properties."

In contrast to a fight-or-flight response, gearing the body up for action, a parasympathetic response slows heart rate and breathing and constricts the pupils.

"These visual cues signal the body to be cautious, while also closing off the body, as if to limit its exposure to something that could be harmful," Ayzenberg says.

The authors theorize that clusters of holes may be evolutionarily indicative of contamination and disease—visual cues for rotten or moldy food or skin marred by an infection.

The subjects involved in the experiments were college students who did

not report having tryphobia. "The fact that we found effects in this population suggests a quite primitive and pervasive visual mechanism underlying an aversion to [holes](#)," Lourenco says.

Since the time of Darwin, scientists have debated the relation between fear and disgust. The current paper adds to the growing evidence that—while the two emotions are on continuums and occasionally overlap—they have distinct neural and physiological underpinnings.

"Our findings not only enhance our understanding of the visual system but also how visual processing may contribute to a range of other phobic reactions," Ayzenberg says.

More information: Vladislav Ayzenberg et al. Pupillometry reveals the physiological underpinnings of the aversion to holes, *PeerJ* (2018). [DOI: 10.7717/peerj.4185](https://doi.org/10.7717/peerj.4185)

Provided by Emory University

Citation: Aversion to holes driven by disgust, not fear, study finds (2018, January 4) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-01-aversion-holes-driven-disgust.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
