

Brain imaging predicts language learning in deaf children

January 15 2018



Credit: CC0 Public Domain

In a new international collaborative study between The Chinese University of Hong Kong and Ann & Robert H. Lurie Children's Hospital of Chicago, researchers created a machine learning algorithm

that uses brain scans to predict language ability in deaf children after they receive a cochlear implant. This study's novel use of artificial intelligence to understand brain structure underlying language development has broad reaching implications for children with developmental challenges. It was published in the *Proceedings of the National Academy of Sciences*.

"The ability to predict [language](#) development is important because it allows clinicians and educators to intervene with therapy to maximize language learning for the child," said co-senior author Patrick C. M. Wong, PhD, a cognitive neuroscientist, professor and director of the Brain and Mind Institute at The Chinese University of Hong Kong. "Since the brain underlies all human ability, the methods we have applied to [children](#) with [hearing](#) loss could have widespread use in predicting function and improving the lives of children with a broad range of disabilities" said Wong.

A cochlear implant is the most effective treatment for children born with significant hearing loss when hearing aids are not enough for the child to develop age appropriate listening and [language ability](#). Decades of research have shown that early cochlear implantation is critical. Although a cochlear implant enables many children with hearing loss to understand and develop speech, some children lag behind their normal hearing peers despite receiving an implant as an infant or toddler. Helping these children achieve the language and literacy of [hearing children](#) is important and the focus of much research, as these skills are critical to academic success, social and emotional well-being and employment opportunities.

"So far, we have not had a reliable way to predict which children are at risk to develop poorer language. Our study is the first to provide clinicians and caregivers with concrete information about how much language improvement can be expected given the child's brain

development immediately before surgery," said co-senior author Nancy M. Young, MD, Medical Director, Audiology and Cochlear Implant Programs at Lurie Children's, a surgeon and professor at Northwestern University Feinberg School of Medicine. "The ability to forecast children at risk is the critical first step to improving their outcome. It will lay the groundwork for future development and testing of customized therapies."

"A one-size-fits-all intensive therapy approach is impractical and may not adequately address the needs of those children most at risk to fall behind," added Wong.

Erin Ingvalson, assistant professor at Florida State University who began work on the project as a post-doctoral student at Northwestern University, said "our goal is to eliminate the gap in language outcomes often found when children with hearing loss are compared to those with normal hearing."

"The ability to optimize therapy for each child with [hearing loss](#) will transform many lives," said Ingvalson.

Successful hearing and spoken language development depends on both the ear and the brain. Hearing loss early in life deprives the auditory areas of the brain of stimulation, which causes abnormal patterns of [brain](#) development.

"We used MRI to capture these abnormal patterns before cochlear implant surgery and constructed a machine-learning algorithm for predicting language development with a relatively high degree of accuracy, specificity and sensitivity," Wong explained. "Although the current algorithm is built for children with hearing impairment, research is being conducted to also predict [language development](#) in other pediatric populations."

More information: Gangyi Feng et al., "Neural preservation underlies speech improvement from auditory deprivation in young cochlear implant recipients," *PNAS* (2018).

www.pnas.org/cgi/doi/10.1073/pnas.1717603115

Provided by Ann & Robert H. Lurie Children's Hospital of Chicago

Citation: Brain imaging predicts language learning in deaf children (2018, January 15) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2018-01-brain-imaging-language-deaf-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.