

Breastfeeding reduces hypertension risk

January 30 2018



Credit: CC0 Public Domain

A study published in the *American Journal of Hypertension* indicates that women who breastfeed more children, and for longer periods of time, are less likely to suffer from hypertension after they reach menopause. This is less true of obese women, however.

Elevated blood pressure is the greatest single risk factor for disease and mortality. Evidence from epidemiologic data has also shown the beneficial effects of breastfeeding on the health of infants and their mothers. It has been well documented that long-term breastfeeding is associated with reduced children's allergies, celiac disease, obesity, and diabetes mellitus. However, the effects of breastfeeding on [maternal health](#) have been little studied compared with the effects on the children.

Several studies consistently found that absent breastfeeding or premature discontinuation was associated with increased risks of diabetes mellitus, dyslipidemia, metabolic syndrome, [coronary heart disease](#), and cardiovascular diseases. However few studies have established a clear relationship between breastfeeding and hypertension.

This study population comprised 3,119 non-smoking [postmenopausal women](#) aged 50 years or older in the 2010-2011 Korea National Health and Nutrition Examination Survey.

More children breastfed and longer duration of breastfeeding were associated with lower risk of hypertension in postmenopausal women, and degree of obesity and [insulin resistance](#) moderated the breastfeeding-hypertension association. In particular, the highest quintile of number of children breastfed (5 to 11) showed a 51% lower risk of hypertension compared with the lowest quintile (0 to 1). The highest quintile of duration of breastfeeding (96 to 324 months) showed a 45% lower risk of [hypertension](#).

Although a broad range of chronic diseases are not associated with breastfeeding, some common mechanisms have been proposed to underlie the relationships between breastfeeding and these diseases. First, maternal metabolism (e.g., fat accumulation and insulin resistance) may be "reset" by breastfeeding after pregnancy, which decreases the risk of obesity-related diseases. Second, oxytocin release stimulated by

breastfeeding may be associated with the decreased risk of these diseases.

"Our findings endorsed the current recommendations for breastfeeding for the benefit of maternal health in mothers' later lives," said the paper's lead researcher, Nam-Kyong Choi.

More information: Breastfeeding and maternal hypertension, *American Journal of Hypertension* (2017). [DOI: 10.1093/ajh/hpx219](https://doi.org/10.1093/ajh/hpx219)

Provided by Oxford University Press

Citation: Breastfeeding reduces hypertension risk (2018, January 30) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-01-breastfeeding-hypertension.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.