

Complications of reconstruction surgery differ for transgender patients

January 31 2018

The risks of penile reconstruction surgery (phalloplasty) appear higher in female-to-male transgender (transmale) patients undergoing gender confirmation surgery, compared to native male (cismale) patients undergoing phalloplasty for other reasons, reports a study in the February issue of *Plastic and Reconstructive Surgery*, the official medical journal of the American Society of Plastic Surgeons (ASPS).

Complications are also less frequent when reconstruction of the urethra (urethroplasty) is done as a single-stage procedure, according to the new research by ASPS Member Surgeon Jeffrey B. Friedrich, MD, of University of Washington, Seattle, and colleagues. "Our results provide useful information for surgeons and patients to consider when discussing options, risks, and expected outcomes of phalloplasty," Dr. Friedrich comments.

Increased Phalloplasty Risks for Transmale Patients and with Staged Urethral Surgery

Phalloplasty has important benefits not only for transmale patients undergoing [gender confirmation surgery](#), but also for cismale patients who have lost their penis due to congenital or acquired disorders (for example, trauma). However, there are few data on the aggregate complications and outcomes of phalloplasty, including possible differences between these two groups of patients.

Dr. Friedrich and colleagues analyzed the findings of 50 previous studies relating to phalloplasty. Nineteen studies reported on phalloplasty in 869 transmale patients, while 31 studies reported on 482 cisfemale patients. The studies used a wide range of surgical approaches to phalloplasty, and assessed varying outcome measures.

The results showed some important differences in outcome between the two groups. The overall rate of complications related to the urethra was 39.4 percent in transmale patients compared to 24.8 percent in cisfemale patients. Rates of complications involving the tissue flaps used for penile reconstruction were also higher in transmale patients: 10.8 versus 8.1 percent.

Outcomes were also affected by the approach used to reconstruct the urethra: 23 studies including 723 patients used a single-step (primary) approach to urethroplasty, while 13 studies including 201 patients used multiple-step (staged) procedures. The rate of tissue flap-related complications was also lower with primary urethroplasty: 8.6 percent, compared to 16.7 percent for staged procedures.

In addition to fewer complications, single-stage urethral reconstruction was associated with improvement in key outcomes, for both transmale and cisfemale patients. Ability to urinate while standing, sexual functioning, and patient satisfaction were all better with primary versus staged urethroplasty.

In both transmale and cisfemale patients, penile reconstruction has important physical and psychosocial benefits. By synthesizing the results of previous studies, the new analysis provides much-needed information on the complications and outcomes of phalloplasty.

"Transmale patients seem to have a higher incidence of complications following phalloplasty than cisfemale patients," according to the authors.

The higher rate of urethral [complications](#) in transmale patients likely reflects the inherent difficulties of reconstruction in this group of patients. The researchers note that the overall patient satisfaction rate was high: about 84 percent across groups.

"Overall, phalloplasty is an effective and efficacious technique, which benefits patients," Dr. Friedrich and colleagues conclude. They highlight the need for further prospective studies to understand the best options available to different groups of patients. They also call for the development of "standardized and reliable" tools for assessing patient-reported outcomes in transgender [patients](#).

More information: Austin C. Remington et al. Outcomes After Phalloplasty, *Plastic and Reconstructive Surgery* (2017). [DOI: 10.1097/PRS.00000000000004061](#)

Provided by Wolters Kluwer Health

Citation: Complications of reconstruction surgery differ for transgender patients (2018, January 31) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-01-complications-reconstruction-surgery-differ-transgender.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--