

Diabetes management improved in high-risk population through community program

January 31 2018



Nadia S. Islam, PhD, associate professor, Department of Population Health at NYU Langone Health, led a study that examined the impact of community-based health workers in helping an at-risk population better manage diabetes. Credit: NYU Langone Health

An ethnic population at high risk for Type 2 diabetes achieved significant control of the disease through participation in community-based health programs, demonstrating that active intervention and

culturally-sensitive education can reverse the course of certain illnesses.

This according to a randomized controlled trial published January 31 by researchers at NYU School of Medicine's Department of Population Health in the journal *Clinical Diabetes*.

The Diabetes Research, Education, and Action for Minorities, or DREAM trial, took place in several neighborhoods throughout New York City heavily populated by Bangladeshi-Americans, who have a notably higher rate of Type 2 diabetes than the general U.S. population. The study participants were randomly assigned to either a control group or a group that participated in several education sessions with community health workers.

After six months, over 55% of those in the [intervention group](#) had blood sugar levels of seven percent or less—a measure of Type 2 diabetes management also known as Hemoglobin A1c levels—compared to 42% in the control group.

"This is the first study of a community-based program to help New York City's Bangladeshi community control diabetes," says Nadia S. Islam, PhD, principal investigator of DREAM and associate professor of Population Health at NYU School of Medicine. "To the broader issue of disease management, these results add to a growing body of evidence showing that community health worker programs are a low cost and effective strategy for controlling certain chronic diseases."

Bangladeshi immigrants make up the fastest growing segment of South Asians in the U.S., numbering over 3.4 million—with a majority living in New York City. Foreign-born South Asians also are at greater risk of diabetes than the general population. A recent study in New York City concluded that 35% of South Asians had Type 2 diabetes while another study in California found a prevalence of 29%—compared to a rate of

slightly more than 9% among the general U.S. [population](#).

The researchers attribute this unusually high incidence of Type 2 diabetes to several factors: heredity, economic hardship, language barriers, and illiteracy and culture differences. These, in turn, create barriers to adequate healthcare.

To help overcome these barriers, the researchers developed [intervention](#) and health education programs led by community-based health workers, who were hired and trained by centers like NYU Langone, and who often come from the same ethnic background and communities as the populations they serve.

"They understand specific challenges and tailor interventions appropriately, facilitating a community's involvement in its own care," said Chau Trinh-Shevrin, DrPH, associate professor of Population Health at NYU School of Medicine and the study's senior investigator. "Community-based health workers build on existing bonds and encourage leadership around health."

How the Trial Was Conducted

The DREAM study enrolled 336 participants who identified as Bangladeshi; resided in New York City; were between 21 and 75 years old; and had a confirmed diagnosis of Type 2 diabetes—defined as a hemoglobin A1c level of 6.5 percent or higher.

The researchers randomly assigned 176 participants into an intervention group that participated in five, two-hour monthly educational sessions led by bilingual Bangladeshi [health workers](#), covering topics including nutrition and healthy eating; physical activity; complications of Type 2 diabetes; preventive self-care; and stress management and family support. Each individual also participated in two, one-on-one visits

where they set individual health goals.

The other 160 participants were assigned to a control group, who were invited only to the first educational session, an overview of diabetes.

At end of the trial period, 36.3% of the intervention group achieved control or stabilization of their hemoglobin A1c levels (defined as under seven percent) compared with 24.6% of the control group. In addition, 55.2% of intervention group also had decreased A1c levels at the end of the study, compared to 42.5% in the [control group](#).

The researchers also found that both the intervention and [control](#) groups reported overall changes in behavior, including increased [diabetes](#) knowledge, increased weekly physical activity, and weekly monitoring of blood sugar. "This suggests that even minimal community engagement around [health](#) made an impact," Islam says.

Provided by NYU Langone Health

Citation: Diabetes management improved in high-risk population through community program (2018, January 31) retrieved 20 April 2024 from

<https://medicalxpress.com/news/2018-01-diabetes-high-risk-population.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--