


Dietary sugar linked to increasing bacterial epidemics

January 3 2018



This photograph depicts *Clostridium difficile* colonies after 48hrs growth on a blood agar plate; Magnified 4.8X. *C. difficile*, an anaerobic gram-positive rod, is the most frequently identified cause of antibiotic-associated diarrhea (AAD). It accounts for approximately 15–25% of all episodes of AAD. Credit: CDC

The increasing frequency and severity of healthcare-associated outbreaks caused by bacterium *Clostridium difficile* have been linked to the widely used food additive trehalose. A team of researchers discovered that in laboratory tests and animal models, trehalose enhances the virulence of epidemic *C. difficile* lineages that predominate in

patient infections. The study appears in the journal *Nature*.

"*C. difficile* infections have always been a problem in hospitals, but during the last 15 years they have become the most common cause of hospital-acquired infections in developed countries," said corresponding author Dr. Robert Britton, professor of molecular virology and microbiology and member of the Alkek Center for Metagenomics and Microbiome Research and the Dan L Duncan Comprehensive Cancer Center at Baylor College of Medicine.

According to the Center for Disease Control and Prevention, *C. difficile* is a major cause of infectious disease-related death in the United States. In 2015, the CDC reported that *C. difficile* caused almost a half-million infections in patients in a year and 29,000 estimated deaths. The bacteria cause life-threatening inflammation of the colon and diarrhea. Patients 65 years and older are at most risk, and most infections occur in people who have received medical care and antibiotics.

"Our group and others have found that *C. difficile* lineages RT027 and RT078 have become dominant more recently around the globe," said first author Dr. James Collins, a postdoctoral associate in the Britton lab. "These lineages have been present in people for years without causing major outbreaks; in the 1980s they were not epidemic or hypervirulent but after the year 2000 they began to predominate and cause major outbreaks. We wanted to know what had helped these lineages become a major health risk."

Diet and bacterial virulence

Resistance to fluoroquinolone antibiotics is likely one of the factors that is helping lineage RT027 cause epidemics.

"However, fluoroquinolone resistance is also a characteristic of other *C.*

difficile lineages that are not epidemic," Collins said. "We searched for other factors that would help RT027 and RT078 increase their virulence."

The researchers investigated what sources of food RT027 and RT078 preferred. They discovered that these lineages can grow on levels of sugar trehalose that are about 1,000 times lower than those needed by other lineages of these bacteria, giving RT027 and RT078 a major advantage. Each lineage is highly efficient at using trehalose and evolved independent mechanisms to utilize this sugar. To connect the ability to metabolize low levels of trehalose with increased disease severity, the researchers worked with a mouse model of *C. difficile* infection.

"Mice received a strain of the RT027 [lineage](#) of *C. difficile* and a diet with or without low trehalose levels," Collins said. "What the mice ate made a difference to the virulence of the infection; mortality was higher in the group consuming trehalose."

Further experiments showed that increased disease severity in the presence of trehalose could not be explained by the mice having higher numbers of bacteria, instead what made the disease more severe was that RT027 produced higher levels of toxins.

These and other experiments provide evidence that dietary trehalose has contributed to the predominance of epidemic *C. difficile* lineages and to their virulence. Because the genetic factors that allow these bacteria to metabolize trehalose and increase the production of toxins were present well before the outbreaks started, the researchers investigated what could have triggered the epidemics.

"In 2000, trehalose was approved as a food additive in the United States for a number of foods from sushi and vegetables to ice cream, and about three years later the reports of outbreaks with these lineages started to

increase," Britton said. "Other factors may also contribute, but we think that trehalose is a key trigger."

"An important contribution of this study is the realization that what we once considered a perfectly safe sugar for human consumption, can have unexpected consequences," Collins said. "Our study suggests that the effect of trehalose in the diet of patients in hospitals with RT027 and RT078 outbreaks should be further investigated."

More information: J. Collins et al, Dietary trehalose enhances virulence of epidemic *Clostridium difficile*, *Nature* (2018). [DOI: 10.1038/nature25178](https://doi.org/10.1038/nature25178)

Provided by Baylor College of Medicine

Citation: Dietary sugar linked to increasing bacterial epidemics (2018, January 3) retrieved 17 May 2024 from <https://medicalxpress.com/news/2018-01-dietary-sugar-linked-bacterial-epidemics.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--