

Does eating more fish protect you from depression?

January 12 2018

According to the world health organisation (WHO), depression is the greatest single cause of disability worldwide. Therefore, understanding mechanisms leading to depression and how to minimise its risks is very important.

Several studies have reported that fish consumption is associated with a reduced risk of [depression](#). However, these studies remain to be controversial due to inconsistencies between some of them, the method that they used to assess depression and because none have established a clear dose that is required for the observed effects on depression.

A group of Japanese researchers has recently demonstrated that moderate fish intake is recommended for preventing depression in aged Japanese [individuals](#).

Yutaka Matsuoka and colleagues from the National Cancer Centre Japan, the University and Toyama and Keio University School of Medicine, investigated the association between [fish consumption](#) and the risk of psychiatrist-diagnosed [major depressive disorder](#) (MDD) in Japan. The study involved 1181 individuals (aged 63-82) for whom dietary information had been available over the course of up to 25 years, because they had taken part in the Japan Public Health Centre-based (JPHC) Prospective Study. Psychiatric assessment was conducted for each of these individuals using both standard questionnaires and assessment by trained psychiatrists.

The authors assessed the active ingredients in fish that are thought to mediate the anti-depression effect which are n-3 polyunsaturated fatty acids (PUFAs) such as eicosapentaenoic acid (EPA) and docosapentaenoic acid (DPA). The study found that the relationship between fish intake and MDD does not follow a simple linear profile, but rather a so-called reverse J-shaped association is observed. A decreased risk of MDD was observed in individuals with a median intake of 111 g/day of fish, 307 mg/day of EPA or 123 mg/day of DPA.

The authors conclude that "emerging and compelling evidence suggests that diet and nutrition are extremely important factors in the high prevalence of depressive disorders, and our findings provide a basis to further examine the effectiveness of [fish](#) and n-3 PUFA intake for the prevention of MDD in both aged individuals and those with a history of major physical illness."

More information: M Kawata et al. Ablation of neuropsin–neuregulin 1 signaling imbalances ErbB4 inhibitory networks and disrupts hippocampal gamma oscillation, *Translational Psychiatry* (2017). [DOI: 10.1038/tp.2017.20](#)

Provided by Keio University

Citation: Does eating more fish protect you from depression? (2018, January 12) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-01-fish-depression.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--