

## Flu may be spread just by breathing, new study shows; coughing and sneezing not required

January 18 2018





A study participant sits in the Gesundheit II machine, which is used to capture and analyze influenza virus in exhaled breath at the University of Maryland School of Public Health. Credit: University of Maryland



It is easier to spread the influenza virus (flu) than previously thought, according to a new University of Maryland-led study released today. People commonly believe that they can catch the flu by exposure to droplets from an infected person's coughs or sneezes or by touching contaminated surfaces. But, new information about flu transmission reveals that we may pass the flu to others just by breathing.

The study "Infectious virus in <u>exhaled breath</u> of symptomatic seasonal influenza cases from a college community," published in the *Proceedings of the National Academy of Sciences*, provides new evidence for the potential importance of <u>airborne transmission</u> because of the large quantities of infectious virus researchers found in the exhaled breath from people suffering from flu.

"We found that <u>flu cases</u> contaminated the air around them with infectious virus just by breathing, without coughing or sneezing," explained Dr. Milton, M.D., MPH, professor of environmental health in the University of Maryland School of Public Health and lead researcher of this study. "People with flu generate infectious aerosols (tiny droplets that stay suspended in the air for a long time) even when they are not coughing, and especially during the first days of illness. So when someone is coming down with influenza, they should go home and not remain in the workplace and infect others."

Researchers from the University of Maryland, San Jose State University, Missouri Western State University and University of California, Berkeley contributed to this study funded by the Centers for Disease Control and Prevention (CDC) and the National Institutes of Health.

Dr. Milton and his research team captured and characterized influenza virus in exhaled breath from 142 confirmed cases of people with



influenza during natural breathing, prompted speech, spontaneous coughing, and sneezing, and assessed the infectivity of naturally occurring influenza aerosols. The participants provided 218 nasopharyngeal swabs and 218 30-minute samples of exhaled breath, spontaneous coughing, and sneezing on the first, second, and third days after the onset of symptoms.

The analysis of the infectious virus recovered from these samples showed that a significant number of flu patients routinely shed infectious virus, not merely detectable RNA, into <u>aerosol</u> particles small enough to present a risk for airborne transmission.

Surprisingly, 11 (48%) of the 23 fine aerosol samples acquired in the absence of coughing had detectable viral RNA and 8 of these 11 contained infectious virus, suggesting that coughing was not necessary for infectious aerosol generation in the fine aerosol droplets. In addition, the few sneezes observed were not associated with greater viral RNA copy numbers in either coarse or fine aerosols, suggesting that sneezing does not make an important contribution to influenza <u>virus</u> shedding in aerosols.

"The study findings suggest that keeping surfaces clean, washing our hands all the time, and avoiding people who are coughing does not provide complete protection from getting the flu," said Sheryl Ehrman, Don Beall Dean of the Charles W. Davidson College of Engineering at San José State University. "Staying home and out of public spaces could make a difference in the spread of the <u>influenza virus</u>."

According to the authors, the findings could be used to improve mathematical models of the risk of airborne influenza transmission from people with symptomatic illness and to develop more effective public health interventions and to control and reduce the impact of <u>influenza</u> epidemics and pandemics. Improvements could be made to ventilation



systems to reduce transmission risk in offices, school classrooms and subway cars, for example. Meanwhile, we can all heed the advice to stay home, if possible, when we are beginning to get sick to prevent even greater numbers of flu cases. And, get vaccinated—it is not perfect but does prevent a significant amount of severe illness.

**More information:** Jing Yan et al. Infectious virus in exhaled breath of symptomatic seasonal influenza cases from a college community, *bioRxiv* (2017). DOI: 10.1101/194985, www.biorxiv.org/content/early/2018/01/03/194985; *PNAS* www.pnas.org/content/early/201 ... /1716561115.abstract

Provided by University of Maryland

Citation: Flu may be spread just by breathing, new study shows; coughing and sneezing not required (2018, January 18) retrieved 6 May 2024 from https://medicalxpress.com/news/2018-01-flu-required.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.